

Progress report

Intestinal malabsorption in the experimental animal

The study of human malabsorption syndromes has fascinated gastroenterologists for many years, but only recently has any understanding of the basic mechanisms underlying these conditions become available. This lack of knowledge was partially due to the inability of investigators to reproduce the same symptoms in the experimental animal. Much progress in 'experimental gastroenterology' has been made in the last few years, which should eventually assist clinical gastroenterologists in their understanding of human diseases. This review, which condenses and updates an earlier article,¹ presents some of the experimental work of the last decade which has helped us in our comprehension of some of the malabsorption syndromes.

Perhaps the most studied malabsorption syndrome in man is coeliac sprue, a disease characterized by a pathological sensitivity to gluten and by the appearance in the intestine of flattened villi, elongated crypts, and an increased cell turnover²⁻⁴. Large numbers of histochemical studies have been performed on biopsy samples from patients with sprue, and a generalized decrease in enzyme activity within the epithelium has been demonstrated.^{2,3,5-8} On the other hand, a specific diffuse reaction for lysosomal hydrolases within the surface cells has also been observed, and it has been suggested that this is the primary cellular change responsible for the characteristic reaction of the tissue.³ For as the surface cells are destroyed by autolysis by the lysosomal hydrolases, the mitotic patterns of the crypts are accelerated in an attempt at compensation for the increased cell loss. This hypothesis received strong support when it was demonstrated that the diffuse activity of lysosomal enzymes in the affected cells could be reduced by corticosteroid administration,⁹ since such drugs are known to be stabilizers of the lysosomal membrane.¹⁰ What is not yet known is how the sensitivity of these patients to gluten is translated at the cellular level by labilization of the lysosomal membrane of the granules of the epithelial cell. Immunological reactions may be involved in this sensitization.^{11,12}

An untreated intestine of patients with coeliac sprue is the site of moderate to severe malabsorption, as has been shown by perfusion studies *in vivo*¹³⁻¹⁵ and by uptake of radioactive substrates by biopsy samples *in vitro*^{16,17}. The reduced absorption is presumably due to a large extent to the reduction in intestinal surface area.

Increased proliferation of the absorptive cells occurs in many other human diseases,^{18,19} though to a lesser extent, and is always accompanied by similar morphological and cell kinetic patterns;¹⁹ only quantitative differences are observed. It may not be necessary for labilization of the lysosomal membrane to be the trigger responsible for provoking this response; any action that causes the destruction of surface cells and therefore an attempt by intact crypts to compensate for the attack should produce the same overall result.

This situation is clearly observed in three experimental models. The first involves the response of the intestine to the infusion of acid into the lumen, a condition where a preferential attack on the surface cells occurs, together with a resultant compensatory increase in cell turnover in the crypts.²⁰⁻²² Riecken *et al* employed high concentrations of lactic acid for their experiments and observed a typical hyperregenerative mucosal lesion in the rat intestine.²² Unfortunately, extremely severe experimental conditions were required to elicit a typical response, a fact that reduces the usefulness of the experimental model. Nevertheless, these investigators²² were able to distinguish two stages of the hyperregenerative response; first, accelerated epithelial desquamation led to increased mitosis, prolonged crypts, and lengthened villi; then if the lactic acid concentration was further increased, the corrosive action was greater, the crypts were further enlarged, but in this case shortened villi were revealed.

The second experimental condition concerns transmissible gastroenteritis of swine. This is a viral disease in which the organism specifically attacks the surface epithelial cells without affecting the crypts.²³⁻²⁵ As a result, a typical sprue-like syndrome develops, with blunt villi,^{23,25-27} increased mitosis,²⁵⁻²⁷ reduced enzyme activities,^{25,26,28} and malabsorption.^{23,28}

Furthermore, coeliac sprue has been closely mimicked in the experimental animal following injection of triparanol.^{29,30} This drug appears to provoke labilization of the lysosomal membrane within the epithelial cell, liberating acid hydrolases which destroy the surface cells; eventually, a syndrome is induced which is formally identical with human sprue. In addition rats poisoned with triparanol also respond to a gluten-free diet although the normal rat is insensitive to the peptide.³¹ This extremely important discovery indicates that the response of the intestine to a gluten-free diet is somewhat non-specific in nature. But it may possibly be related to the initial reaction provoking the hyperregenerative response of the gut, namely, the labilization of the lysosomal membrane which occurs both in sprue and in triparanol intoxication, though presumably not in porcine transmissible gastroenteritis.

One of the clinical situations in which coeliac-like symptoms arise is kwashiorkor, or infantile protein malnutrition.^{18,32} In this case, no response to a gluten-free diet is observed, and the cause of the morphological changes observed at the cellular level is unknown. The histological alterations are accompanied by malabsorption,^{33,34} reduced enzyme levels,³⁵ and a slight decrease in mitotic index.³² In a recent paper,³⁶ amino-acid uptake *in vitro* by biopsy samples was determined, and it was concluded that the tissues were normal since their transport capacity did not change on treatment. Since no real controls were performed on tissue from normal subjects, this conclusion may be unjustified. On the other hand, rats treated with low-protein diets have been shown to have normal amino-acid absorption,³⁷ despite other changes which indicate the induction of kwashiorkor in the animals.

Marasmus, or malnutrition resulting from caloric deprivation, is a clinical entity which differs considerably from kwashiorkor.^{38,39} At the intestinal level, coeliac-like symptoms are not present, disaccharidase activities are normal,⁴⁰ but there is a thinning of the mucosa and a very strong inhibition of mitosis.^{32,38} Similarities with infantile marasmus have been observed in an experimental model using toads which have been kept without food for several months.⁴¹ The animals were of course emaciated, but the intestine was one of the organs that was preferentially subject to autophagy. There was a strong

reduction in mitotic activity, as indeed has been found in other studies on the effect of starvation in mammals,^{38,42-44} and histological study revealed marked atrophy of the villous structure. Amino-acid transport *in vitro* was reduced, but this could be entirely attributed to the loss of mucosal surface tissue.⁴¹ Similar, but less extreme, atrophy of the mucosa has been observed in blind loops of rat intestine, where mitotic inhibition and malabsorption also occurred.⁴⁵ It has been speculated that this type of villous atrophy is due to a diminution in luminal nutrition,⁴⁶⁻⁴⁸ a factor that might also be responsible for hypertrophy of the intestine following resection or hyperphagia.⁴⁷ (For further discussion on intestinal compensation in response to starvation,^{48,49} intermittent starvation with hyperphagia,⁵⁰ resection,^{47,51} or segmental juxtaposition,^{51,52} all of which may ultimately depend on luminal nutrition, the reader is referred to the reviews cited, particularly that of Dowling⁴⁷ where the relationship of cell turnover and functional maturity to intestinal function in cases of compensation is discussed in detail.) Incidentally, Altmann has recently questioned the importance of luminal nutrition in determining villous size, and has proposed, with the aid of elegant experimental data, that pancreatic and biliary secretions contain a factor responsible for enlargement of the villi.⁵³

One further point concerning intestinal atrophy in undernutrition is of interest. Autophagy of the intestine for nutritional purposes must presumably be delicately controlled, though how this process proceeds is unknown. A hint has been afforded by the demonstration of an increase in lysosomal enzymes in various tissues,^{54,55} including the intestine,⁵⁶ during starvation. It can thus be mooted that this increase can be utilized to break down the tissue for other uses.

The effect of mitotic inhibition *per se* has been the object of numerous studies both in the clinic and in the laboratory animal. The immediate response of the intestine is negligible since the surface cells are not attacked. When they are desquamated at the end of their life cycle, they are no longer replaced and a highly inflammatory bowel results. This situation occurs in the clinic as a result of excessive radiation or after treatment with antimetabolic drugs. Generally, structural changes are observed, but clinically, gastrointestinal symptoms are slight.⁵⁷⁻⁵⁹ This finding has been confirmed in the experimental animal when low doses of radiation or antimetabolics, such as folic acid antagonists or colchicine, have been applied. After a relatively small dose of 5-fluoro-uracil, Roche *et al* have observed a good correlation between glucose transport and the villous cell population,⁶⁰ and Perris demonstrated that after a low dose of irradiation, some transport mechanisms were more sensitive to the intoxication than others,⁶¹ indicating that perhaps the biochemical mechanisms *per se* were sensitive to the radiation.

Often, more vigorous conditions have been employed in the experimental animal, both in the case of irradiation⁶²⁻⁶⁸ and in the case of administration of antimetabolic drugs,⁶⁹⁻⁷⁶ and similar results have been obtained with both models, although the effects of ionizing radiation may be complicated by other cellular effects or by extraintestinal actions.¹ Thus transport of glucose *in vivo* is reduced six hours after irradiation,^{62-65,67,68} probably due to an effect on the adrenals,⁶⁵ before any intestinal damage has been observed. The transport capacity is either normal or slightly above normal at 24 hours, both after irradiation^{63,64,66,67} or administration of antimetabolics.⁷⁴ Then 48 hours after either treatment transport was found to fall off abruptly, according to

both determinations *in vivo*^{66,75} and *in vitro*^{67,68,71,74} and histochemically determined enzymes were concomitantly reduced;^{77,79} this corresponds to the time at which all attacked cells have been desquamated. Certain investigators have noticed that the extremely distorted tissue at this stage is still able to concentrate nutrients to a surprisingly good extent;^{68,74,75} others have noted that normal cells are often interspersed with pathological ones as a result of radiological damage^{78,79} During the recovery stage from either irradiation⁸⁰ or administration of antimetabolic drugs,⁷⁴ there is often a delay in functional normalization despite apparent morphological recovery. This could be ascribed to a temporary increase in cell turnover with a resultant immature cell population during recovery.⁸¹

Particularly in the case of colchicine, histochemical studies have shown that administration of low doses of the drug influence specific enzyme systems of the intestinal mucosa.⁸²⁻⁸⁴ Thus disaccharidases^{83,84} and certain dehydrogenases⁸² are particularly sensitive to colchicine. These findings led Herbst *et al* to propose that colchicine acts upon the mechanism responsible for cell differentiation which is concerned with disaccharidase synthesis, and therefore that the use of small doses of colchicine may provide a useful model for the study of clinical disaccharide intolerance.⁸⁴

There is some evidence that radiation and antimetabolic therapy have different actions on the gut. Hampton has compared and contrasted the structural changes produced by colchicine and by irradiation.⁸⁵ It is possible that radiation not only inhibits mitosis, but also has another effect—one possibility that has been discussed elsewhere¹—is that it labilizes the lysosomal membrane.

If the labilization of the lysosomal membrane may have played a role in some of the other syndromes that have been discussed, it certainly has a primary role in the intestinal response to ischaemia and shock. Severe ischaemia of the small intestine is a rare but catastrophic occurrence in the clinic.^{86,87} Only rapid surgical intervention is capable of preventing death,^{88,89} and even if embolectomy and arterial anastomosis succeed in preventing a fatal outcome,⁸⁸ malabsorption often continues for some time after revascularization has been effected.⁹⁰ Reduced vascularization of intestinal loops due to stenosis of smaller vessels gives rise to a syndrome known as 'intestinal angina' which is less drastic, but which is also often accompanied by malabsorption and may require surgical intervention.^{86,87}

Several experimental studies have been performed to gauge the sequence of events in intestinal ischaemia and to reproduce malabsorption during the recovery stage. Acute ischaemia of a loop of intestine is a dramatic process, damage of the surface cells being evident after only 10 minutes, and complete denudation of the villi being accomplished within half an hour in small rodents.^{91,92} This is accompanied by loss of all absorptive functions.^{92,93} Some authors have used subtotal conditions (such as leaving the collateral circulation intact) and considerably less damage has been observed.^{94,95} Lysosomal enzymes are massively released into the blood stream and lumen on total ischaemia of the gut.⁹⁶⁻⁹⁸ On the other hand, the crypt cells are much more resistant, and in the rat, only start to be destroyed after two hours' total ischaemia.⁹⁷ Just as acute total ischaemia provokes a dramatic destruction, the recovery from these lesions is equally spectacular; it has recently been shown in dogs that mucosae that have been completely destroyed by one hour's ischaemia regain normal structure, function, and microcirculation 24 hours later.⁹⁸ If the crypt cells have been destroyed, an abnormal recovery

pattern would have developed,⁹⁹ as has also been observed in some cases of partial ischaemia.^{92,100} On the other hand, rats surviving one hour's clamping of the superior mesenteric artery appeared to have regained normal villous structure after 48 hours but showed signs of an important malabsorption syndrome.¹⁰¹

The response to shock, both in human subjects and in experimental animals, parallels these findings in intestinal ischaemia.¹⁰² The sensitivity of the intestinal epithelium to hypoxia leads to lysosomal release as a result of the reduced blood flow occurring in shock, and hence ischaemic changes in the mucosa occur.¹⁰³ The primary involvement of the intestine in shock has been elegantly demonstrated in experiments in dogs by Bounous and his coworkers.¹⁰³⁻¹⁰⁵

To the clinician, acute ischaemia of the colon is of greater interest than ischaemia of the small intestine, since it is less rare and since it can also appear as a complication after vascular surgery.^{106,107} As a disease, ischaemic colitis has received considerable attention in recent years,^{108,109} but few attempts have been made at reproducing the syndrome in the experimental animal. Four groups¹⁰⁹⁻¹¹⁴ have induced acute colonic ischaemia in dogs, and have succeeded in reproducing the morphological changes that have been observed in the clinic: total ischaemia of several hours' duration leads eventually to gangrenous destruction of the bowel, whereas shorter periods of acute ischaemia may produce stenosis of the bowel, or the colon may recover its structure completely.^{112,114} The colon is less sensitive to total acute ischaemia than the small intestine, little morphological damage being observed after one hour's total ischaemia.^{113,114} Nevertheless, the discrete changes observed were sufficient to abolish all transport functions.^{113,114} After three hours' ischaemia, the damage was so great that only a proportion of the animals were able to regain structure and function of the ischaemic loop.^{109,114}

Some of the traumas to which the intestine has been subjected, which have been discussed in the foregoing paragraphs, have led to denudation of the intestinal wall and loss of mucosal covering. One problem that has seldom been studied is how the intestine recovers so rapidly from such a disastrous state. Only in the case of colonic ischaemia have cases been recorded where the mutilated area remains free of epithelial covering;^{109,114} the small intestine apparently always recovers, even if the cells remain abnormal. Recently a model has been introduced specifically to study the mode of regeneration of the epithelial covering, namely, preferential destruction of the mucosa by injection of formol into the lumen *in vivo*.¹¹⁵ This investigation, which reveals progression of healthy tissue over the underlying muscular layers until the cells meet in the middle of the damaged tissue, may provide a key for mucosal regeneration.

I am grateful to Professor E. O. Riecken (Marburg-an-der-Lahn) for permitting me to quote work of his that is in press.

J. W. L. ROBINSON
Département de Chirurgie Expérimentale
Hôpital Cantonal Universitaire, Lausanne, Suisse

References

- ¹Robinson, J. W. L. (1972). Experimental intestinal malabsorption states and their relation to clinical syndromes. *Klin. Wschr.*, **50**, 173-185.

- ¹Padykula, H. A., Strauss, E. W., Ladman, A. J., and Gardner, F. H. (1961). A morphologic and histochemical analysis of the human jejunal epithelium in nontropical sprue. *Gastroenterology*, **40**, 735-765.
- ²Riecken, E. O., Stewart, J. S., Booth, C. C., and Pearce, A. G. E. (1966). A histochemical study on the role of lysosomal enzymes in idiopathic steatorrhoea before and during a gluten-free diet. *Gut*, **7**, 317-332.
- ³Riecken, E. O. (1970). Die normale Dünndarmschleimhaut und ihre Veränderungen bei einheimischer Sprue: morphologische und histochemische Befunde. *Dtsch. med. Wschr.*, **95**, 2295-2303.
- ⁴Rubin, W., Ross, L. L., Slesinger, M. H., and Weser, E. (1966). An electron microscopic study of adult celiac disease. *Lab. Invest.*, **15**, 1720-1747.
- ⁵Jos, J., Frézal, J., Rey, J., and Lamy, M. (1967). Étude histochemique de la muqueuse intestinale chez l'enfant. II. La maladie coeliaque. *Ann. Histochim.*, **12**, 165-179.
- ⁶Lojda, Z., Fric, P., Jodl, J., and Chmelik, V. (1970). Cyto-chemistry of the human jejunal mucosa in the norm and in malabsorption syndrome. *Curr. Top. Path.*, **52**, 1-63.
- ⁷Riecken, E. O. (1970). Histochemistry of the small intestine in various malabsorption states. In *Modern Trends in Gastroenterology*, edited by W. I. Card and B. Creamer, **4**, pp. 20-41. Butterworths, London.
- ⁸Wall, A. J., Douglas, A. P., Booth, C. C., and Pearce, A. G. E. (1970). Response of the jejunal mucosa in adult coeliac disease to oral prednisolone. *Gut*, **11**, 7-14.
- ⁹Weissman, G., and Dingle, J. (1961). Release of lysosomal protease by ultraviolet irradiation and inhibition by hydrocortisone. *Exp. Cell Res.*, **25**, 207-210.
- ¹⁰Booth, C. C. (1970). Enterocyte in coeliac disease. *Brit. med. J.*, **3**, 725-731, and **4**, 14-17.
- ¹¹Otto, H. F., and Martin, W. (1971). Zur cytologischen Ultra-struktur des Schleimhautstroma bei Enteropathien, insbesondere bei der idiopathischen Steatorrhoe. *Virchows Arch. path. Anat.*, **353**, 191-206.
- ¹²Holdsworth, C. D., and Dawson, A. M. (1965). Glucose and fructose absorption in idiopathic steatorrhoea. *Gut*, **6**, 387-391.
- ¹³Stewart, J. S., Pollock, D. J., Hoffbrand, A. V., Mollin, D. L., and Booth, C. C. (1967). A study of proximal and distal intestinal structure and absorptive function in idiopathic steatorrhoea. *Quart. J. Med.*, **36**, 425-444.
- ¹⁴Russell, R. I., Allan, J. G., Gerskowitch, V. P., and Robertson, J. W. K. (1972). A study by perfusion techniques of the absorption abnormalities in the jejunum in adult coeliac disease. *Clin. Sci.*, **42**, 735-741.
- ¹⁵Semenza, G., Bircher, J., Mülhaupt, E., Koide, T., Pfenninger, E., Marthaler, Th., Gmünder, U., and Hammerli, U. P. (1969). Arbutin absorption in human small intestine: a simple procedure for the determination of active sugar uptake in peroral biopsy samples. *Clin. chim. Acta*, **25**, 213-219.
- ¹⁶Desjeux, J. F., Sandler, L., Sassier, P., and Lestrade, H. (1971). Acquired and congenital disorders of intestinal transport of D-glucose in children. *Rev. europ. Étud. clin. biol.*, **16**, 364-366.
- ¹⁷Rubin, C. E., Eidelman, S., and Weinstein, W. M. (1970). Sprue by any other name. *Gastroenterology*, **58**, 409-413.
- ¹⁸Riecken, E. O. (1970). Vegeichende biopsische Untersuchungen zur formalen Pathogenese der Dünndarmschleimhautveränderungen bei der einheimischen Sprue. I. Lupenmikroskopische, histologische, morphometrische und zellkinetische Befunde. *Klin. Wschr.*, **48**, 1216-1225.
- ¹⁹Eder, M. (1965). Experimentelle Untersuchungen über Schädigungen der Darmschleimhaut. *Verh. dtsch. Ges. Path.*, **49**, 330-333.
- ²⁰Hamilton, J. R. (1967). Prolonged infusion of the small intestine of the rat—effect of dilute solutions of lactic acid on fat absorption and mucosal morphology. *Pediat. Res.*, **1**, 341-353.
- ²¹Riecken, E. O., Bloch, R., Menge, H., Idelberger, K., Kramer, F., Miller, B., and Lorenz-Meyer, H. (1972). Morphologische und funktionelle Befunde nach Milchsäuredauerinfusion in das Jejunum der Ratte. *Z. Zellforsch.*, in press.
- ²²Haeltermann, E. O. (1972). On the pathogenesis of transmissible gastroenteritis of swine. *J. Amer. vet. med. Ass.*, **160**, 534-540.
- ²³Pensaert, M., Haeltermann, E. O. and Burnstein, T. (1970). Transmissible gastroenteritis of swine: virus-intestinal cell interactions. I. Immunofluorescence, histopathology and virus production in the small intestine through the course of infection. *Arch. ges. Virusforsch.*, **31**, 321-334.
- ²⁴Thake, D. C. (1968). Jejunal epithelium in transmissible gastroenteritis of swine: an electron microscopic and histochemical study. *Amer. J. Path.*, **53**, 149-168.
- ²⁵Maronpot, R. R., and Whitehair, C. K. (1967). Experimental sprue-like small intestinal lesions in pigs. *Canad. J. comp. Med.*, **31**, 309-316.
- ²⁶Hooper, B. E., and Haeltermann, E. O. (1969). Lesions of the gastrointestinal tract of pigs infected with transmissible gastroenteritis. *Canad. J. comp. Med.*, **33**, 29-36.
- ²⁷Kelly, M., Butler, D. G., and Hamilton, J. R. (1972). Transmissible gastroenteritis in piglets: A model of infantile viral diarrhea. *J. Pediat.*, **80**, 925-931.
- ²⁸McPherson, J. R., and Shorter, R. G. (1965). Intestinal lesions associated with triparanol: a clinical and experimental study. *Amer. J. dig. Dis.*, **10**, 1024-1033.
- ²⁹Riecken, E. O., Rosenbaum, R., Bloch, R., Menge, H., Ritt, E., Aslan, M., and Dölle, W. (1969). Tierexperimentelle Untersuchungen zur Frage der Spezifität der Dünndarmschleimhautveränderungen bei der einheimischen Sprue. *Klin. Wschr.*, **47**, 202-214.
- ³⁰Bloch, R., Menge, H., Martini, G. A., and Riecken, E. O. (1972). Effect of gluten on glucose absorption from the small intestine in experimentally induced malabsorption in rats. *Digestion*, **7**, 139-146.
- ³¹Brunser, O., Reid, A., Monckeberg, F., Maccioni, A., and Contreras, I. (1968). Jejunal mucosa in infant malnutrition. *Amer. J. clin. Nutr.*, **21**, 976-983.
- ³²Bowie, M. D., Barbezat, G. O., and Hansen, J. D. L. (1967). Carbohydrate absorption in malnourished children. *Amer. J. clin. Nutr.*, **20**, 89-97.
- ³³James, W. P. T. (1970). Suralg absorption and intestinal motility in children when malnourished and after treatment. *Clin. Sci.*, **39**, 305-318.
- ³⁴Barbezat, G. O., Bowie, M. D., Kaschula, R. O. C., and Hansen, J. D. L. (1967). Studies on the small intestinal mucosa of children with protein-calorie malnutrition. *S. Afr. med. J.*, **41**, 1031-1036.
- ³⁵Woodd-Walker, R. B., Hansen, J. D. L., and Saunders, S. J. (1972). The in vitro uptake of lysine and alanine by human jejunal mucosa in protein-calorie malnutrition, in gastro-enteritis and after neomycin. *Acta paediat. scand.*, **61**, 140-144.
- ³⁶Kirsch, R. E., Saunders, S. J., and Brook, J. F. (1968). Amino acid transport in experimental protein-calorie malnutrition. *Amer. J. clin. Nutr.*, **21**, 1302-1305.
- ³⁷Monckeberg, B., F. (1966). Alteraciones bioquímicas en la desnutrición infantil. *Nutr. bromatol. toxicol.*, **5**, 31-42.

- ³⁹Whitehead, R. G. (1969). Factors which may affect the bio-chemical response to protein-calorie malnutrition. In *Protein-Calorie Malnutrition*, edited by A. von Muralt, pp. 38-47. Springer, Berlin, Heidelberg, New York.
- ⁴⁰James, W. P. T. (1971). Jejunal disaccharidase activities in children with marasmus and with kwashiorkor. *Arch. Dis. Childh.*, **46**, 218-220.
- ⁴¹Pérez-González, M., and Robinson, J. W. L. (1971). The response of the toad intestine to prolonged starvation: A morphological, functional and metabolic study. *Europ. J. clin. Invest.*, **2**, 15-22.
- ⁴²Brown, H. O., Levine, M. L., and Lipkin, M. (1963). Inhibition of intestinal epithelial cell renewal and migration induced by starvation. *Amer. J. Physiol.*, **205**, 868-872.
- ⁴³Wiebecke, B., Heybowitz, R., Löhns, U., and Eder, M. (1969). Der Einfluss des Hungers auf die Proliferationskinetik der Dünn- und Dickdarmschleimhaut der Maus. *Virchows Arch. Abt. B. Zellpath.*, **4**, 164-175.
- ⁴⁴Altmann, G. G. (1972). Influence of starvation and refeeding on mucosal size and epithelial renewal in the rat small intestine. *Amer. J. Anat.*, **133**, 391-400.
- ⁴⁵Menge, H., Bloch, R., Schaumlöffel, E., and Riecken, E. O. (1970) Transportstudien, morphologische, morphometrische und histochemische Untersuchungen zum Verhalten der Dünndarmschleimhaut im operativ ausgeschalteten Jejunalschnitt der Ratte. *Z. ges. exp. Med.*, **153**, 74-90.
- ⁴⁶Dowling, R. H. (1967). Compensatory changes in intestinal absorption. *Brit. med. Bull.*, **23**, 275-278.
- ⁴⁷Dowling, R. H. (1970). Small bowel resection and bypass: recent developments and effects. In *Modern Trends in Gastroenterology*, edited by W. I. Card and B. Creamer, **4**, pp. 73-104. Butterworths, London.
- ⁴⁸Levin, R. J. (1969). The effects of hormones on the absorptive, metabolic and digestive functions of the small intestine. *J. Endocr.*, **45**, 315-348.
- ⁴⁹Wiseman, G. (1971). Effect of undernutrition on intestinal active transport of sugars and amino acids. In *Metabolic Processes in the Fetus and Newborn Infant* (Nutricia Symposium), edited by J. H. P. Jonxis, H. K. A. Visser, and J. A. Troelstra, pp. 243-269. Stenfort Kroese NV, Leyden.
- ⁵⁰Fábrý, P., Kujalová, V., and Petrásek, R. (1959). Einige Folgeerscheinungen der funktionellen und morphologischen Adaptation an geänderte Kalorienzufuhr. *Nahrung*, **3**, 642-649.
- ⁵¹Dowling, R. H., and Booth, C. C. (1967). Structural and functional changes following small intestinal resection in the rat. *Clin. Sci.*, **32**, 139-149.
- ⁵²Altmann, G. G., and Leblond, C. P. (1970). Factors influencing villus size in the small intestine of adult rats as revealed by transposition of intestinal segments. *Amer. J. Anat.*, **127**, 15-36.
- ⁵³Altmann, G. G. (1971). Influence of bile and pancreatic secretions on the size of the intestinal villi in the rat. *Amer. J. Anat.*, **132**, 166-177.
- ⁵⁴Desai, I. D. (1969). Regulation of lysosomal enzymes. I. Adaptive changes in enzyme activities during starvation and refeeding. *Canad. J. Biochem.*, **47**, 785-790.
- ⁵⁵Filkins, J. P. (1970). Lysosomes and hepatic regression during fasting. *Amer. J. Physiol.*, **219**, 923-927.
- ⁵⁶Desai, I. D. (1971). Regulation of lysosomal enzymes. II. Reversible adaptation of intestinal acid hydrolases during starvation and realimentation. *Canad. J. Biochem.*, **49**, 170-176.
- ⁵⁷Trier, J. S. (1962). Morphologic alterations induced by methotrexate in the mucosa of human proximal intestine. I. Serial observations by light microscopy. *Gastroenterology*, **42**, 295-305.
- ⁵⁸Trier, J. S., and Browning, T. H. (1966). Morphologic response of the mucosa of human small intestine to X-ray exposure. *J. clin. Invest.*, **45**, 194-204.
- ⁵⁹Tarpila, S. (1971). Morphological and functional response of human small intestine to ionizing radiation. *Scand. J. Gastroent.*, **6** Suppl. 12.
- ⁶⁰Roche, A. C., Bognel, J. C., Bognel, C., and Bernier, J. J. (1970). Correlation between the histological changes and glucose intestinal absorption following a single dose of 5-fluorouracil. *Digestion*, **3**, 195-212.
- ⁶¹Perris, A. D. (1966). Whole-body irradiation and active transport in the rat small intestine in vitro. *Radiat. Res.*, **29**, 597-607.
- ⁶²Farrar, J. T., Small, M. D., Bullard, D., and Ingelfinger, F. J. (1956). Effect of total body irradiation on absorption of sugars from the small intestine. *Amer. J. Physiol.*, **186**, 549-553.
- ⁶³Zehnder, H. (1960). Glukoseresorption aus dem isolierten Jejunum der Ratte nach Röntgenbestrahlung. *Radiol. clin. (Basel)*, **29**, 127-137.
- ⁶⁴Wesemann, I., Kuss, B., Breuer, H., and Parchwitz, H.-K. (1962). Wirkung einer Röntgenbestrahlung auf den Glukose- und Fettsäuretransport im Dünndarm der Ratte. *Strahlentherapie*, **119**, 425-434.
- ⁶⁵Hehl, R., and Wilbrandt, W. (1964). Die Wirkung der Röntgenstrahlung auf die Zuckerresorption. *Helv. physiol. pharmacol. Acta*, **22**, 319-322.
- ⁶⁶Zsebök, Z. B., Jánosy, G., Petrányi, G., Jr., and Kévacs, L. (1966). In vitro-Untersuchung des Absorptionsvermögens des Dünndarms nach supacetalear Röntgenbestrahlung. *Strahlentherapie*, **131**, 122-136.
- ⁶⁷Ponz, F., and Lluch, M. (1967). Effect of X irradiation on the active transport of glucose through the intestine of the rat in vivo. *Rev. esp. Fisiol.*, **23**, 117-125.
- ⁶⁸Finck, W. (1969). Tierexperimentelle Untersuchungen zur Pathogenese des akuten intestinalen Strahlensyndroms. 2. Mitteilung. Die intestinale Absorption nach Teilkörperbestrahlung. *Radiobiol. Radiother.*, **10**, 637-646.
- ⁶⁹Millington, P. F., Finean, J. B., Forbes, O. C., and Frazer, A. C. (1962). Studies on the effects of aminopterin on the small intestine of rats. I. The morphological changes following a single dose of aminopterin. *Exp. Cell Res.*, **28**, 162-178.
- ⁷⁰Rybak, B. J. (1962). Electron microscope studies of intestinal lesions. I. Aminopterin-induced lesions in mice. *Gastroenterology*, **42**, 306-318.
- ⁷¹Antonoli, J. A., Robinson, J. W. L., Fasel, J., and Vannotti, A. (1966). Effet du méthotrexate in vivo sur l'intestin grêle du rat. *Gastroenterologia (Basel)*, **106**, 217-224.
- ⁷²Eder, M., Rostock, H., and Vogel, G. (1966). Die Wirkung von Folsäureantagonisten (Methotrexate) auf die Regeneration der Darmschleimhaut. *Virchows Arch. path. Anat.*, **341**, 164-176.
- ⁷³Levin, R. J. (1966). Effect of colchicine on intestinal function in the rat. *Gut*, **7**, 250-257.
- ⁷⁴Robinson, J. W. L., Antonoli, J. A., and Vannotti, A. (1966). The effect of oral methotrexate on the rat intestine. *Biochem. Pharmacol.*, **15**, 1479-1489.
- ⁷⁵Redgrave, T. G., and Simmonds, W. J. (1967). Effect of aminopterin on the absorption of fat into the lymph of unanesthetized rats. *Gastroenterology*, **52**, 54-66.
- ⁷⁶Levin, R. J. (1968). Anatomical and functional changes of the small intestine induced by 5-fluorouracil. *J. Physiol. (Lond.)*, **197**, 73-74P.
- ⁷⁷Spiro, H. M., and Pearse, A. G. E. (1964). A histochemical analysis of the effect of irradiation on murine duodenal mucosa. *J. Path. Bact.*, **88**, 55-60.

- ⁷⁶Jervis, H. R., Donati, R. M., Stromberg, L. R., and Sprinz, H. (1969). Histochemical investigation of the mucosa of the exteriorized small intestine of the rat exposed to X-irradiation. *Strahlentherapie*, 137, 326-343.
- ⁷⁷Herzer, R., Dennhardt, R., and Merker, H. J. (1970). Morphologische Veränderungen am Dünndarm von Ratten nach Ganzkörperstrahlung bei verschiedenen osmotisch induzierten Zuständen der Resorption und Sekretion (elektronenmikroskopische Untersuchungen). *Fortschr. Röntgenstr.*, 113, 101-112.
- ⁷⁸Prasad, K. N., and Osborne, J. W. (1963). Influence of betamercaptoethylamine on mucosal uptake of ⁵⁹Fe by intestine-irradiated rats. *Int. J. radiat. Biol.*, 7, 245-253.
- ⁷⁹Loehry, C. A., and Creamer, B. (1969). Three-dimensional structure of the rat small intestinal mucosa related to mucosal dynamics. *Gut*, 10, 112-120.
- ⁸⁰Myren, J., Luketic, G. C., Ceballos, R., Sachs, G., and Hirschowitz, B. I. (1966). Effects of colchicine on intestinal mucosal dehydrogenases. I. Histochemical observations. *Amer. J. dig. Dis.*, 11, 394-403.
- ⁸¹Cohen, M. I., and McNamara, H. (1970). The effect of colchicine on guinea pig intestinal enzyme activity. *Amer. J. dig. Dis.*, 15, 247-250.
- ⁸²Herbst, J. J., Hurwitz, R., Sunshine, P., and Kretchmer, N. (1970). Effect of colchicine on intestinal disaccharidases: Correlation with biochemical aspects of cellular renewal. *J. clin. Invest.*, 49, 530-536.
- ⁸³Hampton, J. C. (1966). A comparison of the effects of X-irradiation and colchicine on the intestinal mucosa of the mouse. *Radiat. Res.*, 28, 37-59.
- ⁸⁴Debray, C., and Leymarios, J. (1965). *L'Artère Mésentérique Supérieure*. L'Expansion Scientifique Française, Paris.
- ⁸⁵Williams, L. F., Jr. (1971). Vascular insufficiency of the intestines. *Gastroenterology*, 61, 757-777.
- ⁸⁶Shaw, R. S., and Maynard, E. P. (1958). Acute and chronic thrombosis of the mesenteric arteries associated with malabsorption. *New Engl. J. Med.*, 258, 874-878.
- ⁸⁷Hivet, M., and Bogniel, J.-C. (1964). Cinq observations d'insuffisance artérielle mésentérique. *Arch. Mal. Appar. dig.*, 53, 1105-1110.
- ⁸⁸Joske, R. A., Shamma'a, M. H., and Drummey, G. D. (1958). Intestinal malabsorption following temporary occlusion of superior mesenteric artery. *Amer. J. Med.*, 25, 449-455.
- ⁸⁹Rao, S. N., and Wynn-Williams, A. (1969). A comparison of post-mortem and ischaemic changes in the absorptive cells of the jejunal mucosa in mice. *Proc. Univ. Otago med. Sch.*, 47, 89-91.
- ⁹⁰Robinson, J. W. L., Antoniolli, J.-A., and Mirkovitch, V. (1966). The intestinal response to ischaemia. *Naunyn-Schmiedeberg's Arch. exp. Path. Pharmacol.*, 255, 178-191.
- ⁹¹Guthrie, J. E., and Quastel, J. H. (1956). Absorption of sugars and amino acids from isolated surviving intestine after experimental shock. *Arch. Biochem.*, 62, 485-496.
- ⁹²David, H., and Uerlings, I. (1967). Elektronenmikroskopische Befunde am Dünndarm des Kaninchens nach Gefäßunterbindung. *Exp. Path.*, 1, 30-44.
- ⁹³Chiu, C. J., McArdle, A. H., Brown, R., Scott, H. J., and Gurd, F. N. (1970). Intestinal mucosal lesion in low-flow states. I. A morphological, hemodynamic and metabolic reappraisal. *Arch. Surg.*, 101, 478-483.
- ⁹⁴Bounous, G., and McArdle, A. H. (1969). Release of intestinal enzymes in acute mesenteric ischemia. *J. surg. Res.*, 9, 339-346.
- ⁹⁵Röttger, P., and Oran, M. (1969). Histochemische und biochemische Untersuchungen zur Frage des Enzymgehaltes der Dünndarmschleimhaut nach temporärer Ischämie. *Verh. dtsh. Ges. Path.*, 53, 189-193.
- ⁹⁶Robinson, J. W. L., and Mirkovitch, V. (1972). The recovery of function and microcirculation in small intestinal loops following ischaemia. *Gut*, 13, 784-789.
- ⁹⁷Glotzer, D. J., Villegas, A. H., Anekamaya, S., and Shaw, R. S. (1962). Healing of the intestine in experimental bowel infarction. *Ann. Surg.*, 155, 183-190.
- ⁹⁸Passi, R. B., and Lansing, A. M. (1964). Experimental malabsorption produced by vascular insufficiency. *Canad. J. Surg.*, 7, 332-340.
- ⁹⁹Cameron, G. R., and Khanna, S. D. (1959). Regeneration of the intestinal villi after extensive mucosal infarction. *J. Path., Bact.*, 77, 505-510.
- ¹⁰⁰Messmer, K. (1967). Intestinale Faktoren im Schock: Intestinaler Kreislauf. *Langenbecks Arch. klin. Chir.*, 319, 890-909.
- ¹⁰¹Sutherland, N. G., Bounous, G., and Gurd, F. N. (1968). Role of intestinal mucosal enzymes in the pathogenesis of shock. *J. Trauma*, 8, 350-380.
- ¹⁰²Bounous, G. (1965). Metabolic changes in the intestinal mucosa during hemorrhagic shock. *Canad. J. Surg.*, 8, 332-339.
- ¹⁰³Bounous, G. (1967). Role of the intestinal contents in the pathophysiology of acute intestinal ischemia. *Amer. J. Surg.*, 114, 368-375.
- ¹⁰⁴Bernatz, P. E. (1960). Necrosis of the colon following resection for abdominal aortic aneurysms. *Arch. Surg.*, 81, 373-378.
- ¹⁰⁵Smith, R. F., and Szilagy, D. E. (1960). Ischemia of the colon as a complication in the surgery of the abdominal aorta. *Arch. Surg.*, 80, 806-821.
- ¹⁰⁶Marston, A., Pheils, M. T., Thomas, M. L., and Morson, B. C. (1966). Ischaemic colitis. *Gut*, 7, 1-15.
- ¹⁰⁷Rausis, C. (1972). Contribution expérimentale à l'étude clinique des désordres vasculaires du gros intestin. Thèse de Doctorat, Faculté de Médecine, Université de Lausanne.
- ¹⁰⁸Boley, S. J., Krieger, H., Schultz, L., Robinson, K., Siew, F. P., Allen, A. C., and Schwartz, S. (1965). Experimental aspects of peripheral vascular occlusion of the intestine. *Surg. Gynec. Obstet.*, 121, 789-794.
- ¹⁰⁹De Villiers, D. R. (1966). Ischaemia of the colon: An experimental study. *Brit. J. Surg.*, 53, 497-503.
- ¹¹⁰Marston, A., Marcuson, R. W., Chapman, M., and Arthur, J. F. (1969). Experimental study of devascularization of the colon. *Gut*, 10, 121-130.
- ¹¹¹Rausis, C., Robinson, J. W. L., Mirkovitch, V., and Saegesser, F. (1972). Ischémie colique nécrosante et gangreneuse: Faits cliniques et recherche expérimentale. *Helv. chir. Acta*, 39, 251-258.
- ¹¹²Robinson, J. W. L., Rausis, C., Basset, P., and Mirkovitch, V. (1972). Functional and morphological response of the dog colon to ischaemia. *Gut*, 13, 775-783.
- ¹¹³Mirkovitch, V., Robinson, J. W. L., Fischer, A., and Winistörfer, B. (1972). Régénération de la muqueuse intestinale après nécrose sélective: étude fonctionnelle, morphologique et microcirculatoire. *Helv. chir. Acta*, in press.