

in the treatment of
peptic ulcer

ROTER

has particular
advantages



ROTER achieves a high cure rate and often gives effective control of even refractory chronic cases which have resisted other treatments.

ROTER in many cases obviates the need for hospitalization or surgical intervention.

ROTER helps patients to keep ambulant and symptom-free with a minimum of medication and with few, if any, dietary restrictions.

ROTER in general practice enables the physician to take adequate care of his gastric cases without frequent or prolonged supervision.

ROTER has no known side effects.

ROTER requires no supplementary medication such as sedatives or anti-spasmodics.

ROTER shows a saving in cost when compared with conventional methods of treatment.

REFERENCES

Therapie der Gegenwart, 1953, 3, 108
Gazz. Med. Ital., 1954, 113, 343
Die Medizinische, 1955, 4, 163
British Medical Journal, 1955, 2, 827
C.R.S. Biol., 1956, 150, 1019
Am J. Gastroenterology, 1957, 28, 439
Wiener Med. W., 1958, 108, 218
Ärztliche Praxis, 1958, 10, 766
Journal Lancet, 1960, 80, 37

FORMULA

Each Roter tablet contains:
Bismuth Subnitrate Roter (350 mg.), Magnesium Carbonate (400 mg.), Sodium Bicarbonate (200 mg.) and Cortex Rhamni Frangulae (25 mg.).

PACKINGS OF ROTER TABLETS

Tins of 40 and 120, also dispensing sizes, 360 and 720 (P.T. exempt). Basic N.H.S. price: Under 1/- for 10 tablets.

ROTER TABLETS are in Category "S". Literature on Request

F.A.I.R. LABORATORIES LTD · TWICKENHAM · MIDDLESEX

**NOW
IN**




**JUST
TEAR
OPEN!**

*

a more convenient and economical
presentation of hydrocortisone
for ulcerative colitis

EFCORTELAN SOLUTION

TRADE MARK




Given as a nightly rectal drip, soluble hydrocortisone (Efcortelan) has been demonstrated to afford rapid alleviation of symptoms of ulcerative colitis in most patients. Symptomatic relief is accompanied by an improvement in the sigmoidoscopic findings and in the colonic mucosa.

The method of treatment is quite suitable for the patient to carry out himself provided the attack is not severe.

With soluble Efcortelan, hormonal side-effects are avoided. If further attacks occur they can usually be

reduced by another course of treatment.



Only one of the new Efcortelan Solution Tablets in the appropriate amount of normal saline (100-120 ml) is needed to prepare the retention enema. It is therefore the ideal presentation for home treatment.


EFCORTELAN SOLUTION TABLETS

Foil strip of 7 tablets (one week's treatment). Each tablet contains 100 mg hydrocortisone (as water soluble sodium succinate).

Subsidiary Companies or Agents in most countries

**GLAXO LABORATORIES LTD.
GREENFORD, MIDDLESEX
BYRON 3434**

transid
brand



transid

A unique antacid.....

polyaminostyrene

insoluble, non-alkaline ion-exchange resin

Of the antacids at present available for the treatment of peptic ulcer, 'Transid' most nearly approaches the ideal, defined ("The Pharmacological Basis of Therapeutics", Goodman & Gilman, 1955) as

"that which does not cause systemic alkalosis, does not cause 'rebound' stimulation of the secretion of acid, interferes least with digestive processes, does not cause diarrhoea or constipation, is not irritative to the stomach, does not release carbon dioxide upon reacting with hydrochloric acid and, lastly, is efficient in neutralising gastric juices."

'Transid' is active only against *excess* acid; it also reduces pepsin activity. Otherwise, 'Transid' is completely inert; it does not interfere with digestion or absorption and cannot cause alkalosis, constipation or diarrhoea. Clinical experience suggests that 'Transid' may succeed in cases of peptic ulcer in which medical treatment with orthodox antacids has failed.

INDICATIONS:

Peptic ulcer; gastric hyperacidity from whatever cause; dyspepsia of pregnancy.

No ill-effect can come from too much polyaminostyrene and the correct dose is that required to control the symptoms (initially, usually two to four tablets every two or three hours).

PACK AND PRICE:

100 tablets in foil pack, basic NHS cost, 10s.

May we send you a copy of our Standard Publication on 'Transid'? It contains much more information than can be included in a journal advertisement.

DUNCAN FLOCKHART OF EDINBURGH

The Doctors' House

Duncan, Flockhart & Co Ltd. Wheatfield Road, Edinburgh 11



*“I cannot eat but lytel mete
My stomach is not goode . . .”*

Anon 16th cent.

In diarrhoeic conditions due to infection, clinical experience has shown ‘Sulphamezathine’ to be far superior to the non-absorbable sulphonamides.

‘Sulphamezathine’ can be given with safety

to patients of all ages, and renal complications are most unlikely to occur.*

Furthermore, ‘Sulphamezathine’ dispenses with the need for using antibiotics—especially those of the costly “broad-spectrum” group.

* Brit. Nat. Form. (Stand. Edn.) 1960, p.45.

Sulphamezathine

(Sulphadimidine B.P.)

Trade mark

Basic N.H.S. cost: 1/7½d per 25 tablets (0.5 g.). Suspension (100 ml.) 3/4d.



NULACIN

"The most liberal treatment of gastric acid"

Nulacin tablets, dissolved slowly in the mouth, provide intragastric milk-alkali drip therapy, obviating the necessity for a tube. Treatment is easy and convenient for the ambulant as well as the bedridden patient. Nulacin may be relied on to provide an effective treatment for peptic ulcer as well as other conditions associated with gastric hyperacidity.

No undesirable side effects occur and no cases of alkalosis or acid rebound due to Nulacin have ever been reported. The safety and effectiveness of Nulacin therapy has been proven by 'in vivo' clinical work carried out in many countries.

*Proc. R. Soc. Med., 1958, 51, 1063.


Nulacin tablets have no B.P. equivalent. The basic N.H.S. price of the 25 tablet tube is 2/-.

simply

safely

effectively

economically



NULACIN is a **HORLICKS** pharmaceutical product

Further information is available from:
 HORLICKS LIMITED, PHARMACEUTICAL DIVISION, SLOUGH, BUCKS

When
infection
is at
hand



'Streptotriad'

trade mark

brand

TRI-SULPHONAMIDE/STREPTOMYCIN

against bacillary dysentery

for the treatment of the acute phase
and during the convalescent or chronic stage to prevent
the patient becoming a carrier.

Each 'Streptotriad' tablet and teaspoonful of prepared suspension contains:

Streptomycin (as sulphate) B.P. 65 mg.

Sulphamerazine B.P. 65 mg.

Sulphadiazine B.P. 100 mg.

Sulphathiazole B.P.C. 100 mg.



An M&B brand Medical Product


MANUFACTURED BY **MAY & BAKER LTD**

MA 8256

DISTRIBUTORS PHARMACEUTICAL SPECIALITIES (MAY & BAKER LTD · DAGENHAM

ASPIRIN IS SAFER WHEN IT IS SOLUBLE

*Soluble aspirin is safer
when it is stable*



N.H.S. Basic Price 1 2/6 for 500 tablets in foils.

SOLPRIN provides soluble aspirin in its most stable form and dissolves rapidly in water to form a solution of calcium acetylsalicylate.

SOLPRIN is easy to take and palatable. It has proved rapid and reliable in its analgesic effect and is extremely well tolerated, even in prolonged and massive dosage.


SOLPRIN is not advertised to the public and is available only on prescription. Clinical samples and literature will be supplied on request.

SOLPRIN

REGD.

Soluble and substantially neutral

**GASTRIC ACID
SECRETION
BACK TO NORMAL
WITH**



NACTON

REGD. TRADE MARK

**NACTON GIVES RELIEF FROM PAIN IN
PEPTIC ULCER AND HYPERACIDITY**

- * 50%—75% reduction of gastric acid secretion
- * Activity prolonged for as much as 8 hours
- * Selective action
- * No disturbing side effects

Each Nacton tablet contains 2 mg. of (1-methyl-2-pyrrolidyl) methyl benzilate methylmethosulphate.



NACTON is a product of British Research
at **BEECHAM RESEARCH Laboratories Ltd.**
Brentford, England

For pre-operative intestinal sterilisation

SOFRAMYCIN TABLETS

250 mg. oral tablets


Effective against most organisms especially staphylococci, enterococci, enterobacteria including *Proteus* and particularly *Escherichia coli*.

“In several instances there was no growth of organisms at all in the stool after total dosages of 2 or 3 g.”

“Undesirable side-effects from the drug were not found”
Gut, 1960, 1, 323.




250 mg. Soframycin Oral Tablets
Bottles of 20



London, N.W.10. Telephone: LADbroke 6611

ULCERATIVE COLITIS



In Ulcerative Colitis
extensive clinical
experience confirms

SALAZOPYRIN

BRAND OF SULPHASALAZINE
(5-AMINO-4-ACETYLAMINO-2-PYRIDINE)
(5-AMINO-4-ACETYLAMINO-2-PYRIDINE)

to be a drug of
significance in to-day's
management of this
distressing disease

"The present investigation suggests that
salazopyrin is a useful drug." *Gut* 1, 217 (1960)

Outstanding characteristics of Salazopyrin treatment are

- Prompt remission of symptoms
- Bleeding and frequency of stools sharply reduced
- In many cases rectal mucosa heals within a month
- Abdominal cramp controlled
- Fever subsides, appetite improves
- Can be given safely over long periods of time

SALAZOPYRIN is also a safe and effective product in other forms of colitis and regional enteritis.

IRI PHARMACIA
OF SWEDEN


Division for Sales, 1, West, London, W.1, England

Pro-Banthine

*Relief
of pain
in peptic
ulcer*

*Pro-Banthine plays a part
in relief of pain with Duclalan*

W. H. WEAVER



Using the ENEMA in this day and age?

In the preparation of patients for operation, particularly operations on the abdomen, Dulcolax can effectively replace the enema as a method of bowel clearance. Two tablets in the evening followed by one suppository on the morning of the operation are normally sufficient. Unpleasantness for both patient and nurse is thus avoided.

Dulcolax is equally effective as a treatment for post-operative constipation. Being free from side-effects it can safely be used in pregnancy and for young and old patients alike.

If you are still using the enema, you are invited to compare it with the safety, the efficacy, the convenience of Dulcolax.

DULCOLAX*

(4,4'-diacetoxy-diphenyl)-(pyridyl-2)-methane




Packed and distributed in the U.K. by Pfizer Ltd., Sandwich, Kent for
C. H. BOEHRINGER SOHN, INGELHEIM am RHEIN
Registered proprietors of the Trade Mark

*Regd. Trade Mark

12015

**ALL YOUR
EGGS . . .**



IN ONE BOTTLE

Not really, because Aminosol is a casein hydrolysate. It does, however, provide adequate protein parenterally where oral nutrition is impracticable.

Supplied as Aminosol Glucose Vitrum (*Aminosol G.V.*)

307 calories per litre and

Aminosol Glucose Ethanol Vitrum (*Aminosol G.E.V.*)

662 calories per litre

In 500 and 1000 ml. bottles. Aminosol Glucose Vitrum also provided in 100 ml. paediatric size. Further information on request

AMINOSOL

Vitrum

Is there a **DROXALIN** in the house?



Gastric crises occur even in Doctors' houses—so we learn from letters received. When they do, the cry goes forth "Is there a Droxalin in the house?" Doctor must then perforce produce two Droxalin Tablets from *somewhere*—probably the top drawer in his surgery desk. Whether the trouble be indigestion, Gastric Ulcer or Sickness of Pregnancy "Two Tabs. Droxalin" bring speedy relief. They are also easy to chew and *most* pleasant to taste.

This medical confidence in Droxalin is reflected also in its wide prescribing, hence the ever-growing question in a Gastric Crisis "Is there a Droxalin in the House?"

In the letters we receive doctors speak highly of Droxalin as:

Giving rapid relief to the abdominal pain and discomfort due to Hyperchlorhydria and Peptic Ulcer.

Being an excellent tablet for indigestion, and in particular for the indigestion of pregnancy.

Effective in acute Dyspepsia and accompanying Insomnia. Giving instant relief from abdominal discomforts and Duodenal Ulcer.

ACTIVE INGREDIENTS:

Dried Aluminium Hydroxide Gel. B.P. 2.5 grains.
Magnesium Trisilicate B.P. 2.5 grains.

PACKING AND PRICE

Droxalin Tablets are hygienically packed in film strips of 6, in cartons of 30 and dispensing packs of 300. Prescribable on E.C.10. N.H.S. cost 2/2 for 50 tablets.

DROXALIN
BFGD
TABLETS

ACID ADSORBENT

speed of action plus palatability

PHILLIPS, SCOTT & TURNER - SURBITON - SURREY

M3

M3

STAPHYLOCOCCAL SUPERINFECTIONS

"Patients in hospital who have had an operation, and a course of 'broad-spectrum' antibiotics, sometimes develop diarrhoea which may be severe and even fatal."

Lancet Annotations 25th February, 1961

ENPAC

LACTOBACILLUS ACIDOPHILUS PREPARATION

For the prophylaxis and treatment of staphylococcal super infections.

"It was found that the staphylococci increased in both groups at the beginning of antibiotic treatment. Staphylococci continued to increase in the patients taking the antibiotic alone, but in those taking 'ENPAC' there was a pronounced drop in numbers."

Gordon D., Macrae J., & Wheeler D. M. W. Lancet—May 4th, 1957. pp. 899-901

Samples and literature will be sent to the medical profession on request to:—

WILTS UNITED DAIRIES LIMITED · NUTRITION DEPARTMENT · TROWBRIDGE · WILTS

★ *Some Outstanding Livingstone Publications* ★

SURGERY OF THE ACUTE ABDOMEN

By JOHN A. SHEPHERD, V.R.D., M.D., M.Ch., F.R.C.S.(Ed.), F.R.C.S.(Eng.), with Foreword
by SIR ZACHARY COPE

1238 pages

104 illustrations

£5

PEPTIC ULCERATION A Symposium for Surgeons

By CHARLES WELLS, M.B., F.R.C.S., and JAMES KYLE, M.Ch., F.R.C.S.(Eng.), F.R.C.S.I.

272 pages

69 illustrations

42s.

A TEXTBOOK OF SURGICAL PHYSIOLOGY

By R. A. JAMIESON, M.B., F.R.C.S.(Ed.), and A. W. KAY, M.D., Ch.M., F.R.C.S.(Ed.)

631 pages

233 illustrations

55s.

Two Books by E. S. R. HUGHES, M.D., M.S., F.R.C.S.(Eng.), F.R.A.C.S.

SURGERY OF THE ANUS, ANAL CANAL AND RECTUM **SURGERY OF THE COLON**

316 pages

398 illustrations

50s.

424 pages

339 illustrations

63s.

E. & S. LIVINGSTONE LTD. - TEVIOT PLACE - EDINBURGH