

Ease the spasm. Ease the mind.

LIBRAXIN

clidinium bromide and chlordiazepoxide

Clidinium bromide to calm the gut. Chlordiazepoxide to calm the mind.

Indications For the control of hypersecretion, hypermotility and emotional factors associated with gastro-intestinal disorders, such as nervous dyspepsia, peptic ulcer, cardiospasm, pylorospasm, nervous or irritable colon.

Dosage 1 or 2 tablets three or four times daily. In elderly patients, it is recommended that the initial dose be 1 tablet twice daily.

Contra-indications Because of its anticholinergic effects, Libraxin should not be given to patients suffering from glaucoma or prostatic enlargement.

Precautions Patients should avoid alcohol while under treatment with Libraxin, since the individual



Sauter

response cannot be foreseen. Patients' reactions (driving ability, operation of machinery, etc.) may be modified to a varying extent, depending on dosage and individual susceptibility. The established medical principle of prescribing medicaments in early pregnancy only when absolutely indicated should be observed.

Side-effects Side-effects are infrequent and are controlled by reduction of dosage. They include

drowsiness, muscle weakness, dryness of the mouth, blurring of vision, constipation and hesitancy of micturition.

Presentation Libraxin tablets containing 5mg chlordiazepoxide and 2.5mg clidinium bromide in packings of 100 and 500.

Basic NHS Cost 1 tablet 3 times daily 10.2p/day ex 500 pack.

Licence Number 0031/5024

Licence Holder Sauter Laboratories
Division of Roche Products Limited, PO Box 8
Welwyn Garden City, Hertfordshire AL7 3AY
Libraxin is a trade mark

J486074/284

Have you the abilities?



gives you:

- Viewability
- Cleanability
- Immersibility
- Durability
- Serviceability
- Interchangeability








Only the Olympus Endoscopy System with a range of flexible endoscopes for oesophagoscopy, gastroscopy, duodenoscopy, sigmoidoscopy, colonoscopy and bronchoscopy, gives you more abilities – the abilities **you've asked for**.

Only the Olympus Endoscopy System is backed by KeyMed's unique 12-month entirely unconditional guarantee and commitment to customer service that is second to none.



The ultimate fiberscopes by **OLYMPUS** from **KEY MED**





KEY MED
 Specialised Services to Medicine
 KeyMed House, Stock Road
 Southend-on-Sea, Essex SS2 5QH.
 England. Tel: (0702) 616333 (20 lines)
 Telex: 995283 Telefax: (0702) 65677

Gastrozepin is a selective antimuscarinic agent which provides balanced control of gastric secretion without markedly affecting other peripheral receptor sites. This gastro-selective action means that, in practice, Gastrozepin is a well-tolerated drug which heals peptic ulcers.

Gastrozepin DOES NOT . . .

- rely on acid reduction alone
- rely on pepsin reduction alone
- rely on mucosal protection alone
- profoundly affect intragastric pH

Gastrozepin DOES . . .

- relieve daytime pain
- relieve night-time pain
- reduce antacid intake
- heal peptic ulcers with one 50 mg tablet b.d.

For the treatment of peptic ulcer


Twice daily

GASTRO SELECTIVE


Gastrozepin[®]

pirenzepine




Prescribing Information; Presentation: White tablets each containing 50 mg of pirenzepine dihydrochloride scored on one face with 'G' on one side of the score, and '50' on the other. The obverse is impressed with the symbol . **Uses:** Gastrozepin is indicated in the treatment of gastric and duodenal ulcers. **Dosage:** 50 mg at bedtime and in the morning before meals. In severe cases the total daily dose may be increased to 150 mg in divided doses. Continuous therapy may be recommended for up to three months. **Contra-indications, Warnings etc:** Interaction with sympathomimetics and monoamine oxidase inhibitors and Gastrozepin is a theoretical possibility. Gastrozepin is not recommended during pregnancy although in animal

experiments no teratogenic effects were noted. Breast milk concentration after therapeutic doses is unlikely to affect the infant. **Side effects:** occasionally transitory dry mouth and accommodation difficulty may occur. Treatment of overdose is entirely symptomatic. There is no specific antidote. **Basic NHS price:** 50 mg tablets, 60 £20.50. **Product Licence No.:** 50 mg tablets, PL00314/0260.

 Further information is available on request
The Boots Company PLC, Nottingham

Gastrozepin[®] Trade Mark



Gastro-technology

Tagamet

cimetidine
acid controlled

Prescribing Information. Presentations 'Tagamet' Tablets, PL 0002/0092, each containing 400 mg cimetidine. 56, £16.61. 'Tagamet' Tablets, PL 0002/0063, each containing 200 mg cimetidine. 500, £74.15. 'Tagamet' Syrup, PL 0002/0073, containing 200 mg cimetidine per 5 ml. 500 ml, £20.43. **Indications** Duodenal ulcer, benign gastric ulcer, recurrent and stomal ulceration, oesophageal reflux disease. Other conditions where reduction of gastric acid is beneficial: prophylaxis of stress-induced gastrointestinal haemorrhage and of acid aspiration (Mendelson's) syndrome, malabsorption and fluid loss in short bowel syndrome. Zollinger-Ellison syndrome. **Dosage Usual dosage:** Adults. *Duodenal ulcer*, 400 mg b.d. with breakfast and at bedtime, or 200 mg t.d.s. with meals and 400 mg at bedtime (1.0 g/day) for at least 4 weeks. To prevent relapse, 400 mg at bedtime or 400 mg morning and at bedtime for at least 6 months. *Benign gastric ulcer*, 200 mg t.d.s. with meals and 400 mg at bedtime (1.0 g/day) for at least 6 weeks. *Oesophageal reflux disease*, 400 mg t.d.s. with meals and 400 mg at bedtime (1.6 g/day) for 4 to 8 weeks. *Prophylaxis of stress-induced gastrointestinal haemorrhage*, up to 2 g a day, divided, to maintain intragastric pH above 4. *Prophylaxis of acid aspiration syndrome*, 400 mg 90-120 mins before induction of general anaesthesia. 400 mg at start of labour then 200 mg 2-hourly as necessary, suggested maximum 1.6 g. Do not use 'Tagamet' syrup. *Zollinger-Ellison syndrome*, up to 2 g a day, divided. *Recurrent and stomal ulceration and short bowel syndrome*, 200 mg t.d.s. and 400 mg at bedtime (1.0 g/day). *N.B. For full dosage instructions see Data Sheet.* **Cautions** Impaired renal function: reduce dosage (see Data Sheet). Potentiation of oral anticoagulants, phenytoin and theophylline (see Data Sheet). Prolonged treatment: observe patients periodically. Exclude malignancy in gastric ulcer. Care in patients with compromised bone marrow (see Data Sheet). Avoid during pregnancy and lactation. **Adverse reactions** Diarrhoea, dizziness, rash, tiredness. Rarely, mild gynaecomastia, reversible liver damage, confusional states (usually in the elderly or very ill), interstitial nephritis, acute pancreatitis. **Legal category** POM. 9.12.83.

Smith Kline & French Laboratories Limited, Welwyn Garden City, Hertfordshire AL7 1EY. © 1984 Smith Kline & French Laboratories Limited
'Tagamet' is a trade mark

SK&F

A BETTER CHOICE EVERY TIME

IT WORKS In the treatment of ulcerative colitis, Colifoam is as effective as steroid enemas. At the same time it has been shown that patients find the foam easier to retain.^{1,2}

PATIENTS PREFER IT Colifoam is far more comfortable, more convenient and more acceptable than enemas. Patients also find it easier to administer and that it causes less interference in their daily lives.

IT COSTS LESS Surprisingly, despite the fact that it's just as effective and far more comfortable, Colifoam is less expensive. In fact, it can cost up to 1/3 less per dose than a standard proprietary enema.³



IT'S SAFER

Recent clinical data shows Colifoam has extremely low levels of systemic absorption,⁴ lower than proprietary prednisolone enemas.⁵ Therefore, there is less potential for adrenal suppression which means that Colifoam may be considered safer in long-term use.

COLIFOAM

hydrocortisone acetate foam


IN DISTAL INFLAMMATORY BOWEL DISEASE. A BETTER CHOICE EVERY TIME.

Presentation White odourless aerosol foam containing hydrocortisone acetate PhEur 10%. **Uses** Anti-inflammatory corticosteroid therapy for the topical treatment of ulcerative colitis, proctosigmoiditis and granular proctitis. **Dosage and administration** One applicatorful inserted into the rectum once or twice daily for two or three weeks and every second day thereafter. Shake can vigorously before use (illustrated instructions are enclosed with every pack). Satisfactory response usually occurs within five to seven days. **Contra-indications, warnings etc.** Local contra-indications to the use of intrarectal steroids include obstruction, abscess, perforation, peritonitis, fresh intestinal anastomoses and extensive fistulae. General precautions common to all corticosteroid therapy should be observed during treatment with Colifoam. Treatment should be administered with caution in patients with severe ulcerative disease because of their predisposition to perforation of the bowel wall. Safety during pregnancy has not been fully established. **Pharmaceutical precautions** Pressurized container. Protect from sunlight and do not expose to temperatures above 50°C. Do not pierce or burn even after use. Do not refrigerate. Shake vigorously before use. Keep out of reach of children. For external use only. **Legal category** POM. **Package quantities** Aerosol canister containing 25g (approx. 14 applications). **Basic NHS cost** 25g plus applicator, £7.40. **Further Information** One applicatorful of Colifoam provides a dose of approximately 125mg of hydrocortisone acetate, similar to that used in a retention enema, for the treatment of ulcerative colitis, sigmoiditis and proctitis. **Product Licence No.** 0036/0021. **References** 1. Ruddell W/SJ, et al. Gut 1980; 21: 885-889. 2. O'Donoghue D. Modern Medicine, December 1981; 45. 3. Source: Mims. 4. Barr W/H, Kline B, Beightol L, Ziass A. Medical College of Virginia/Virginia Commonwealth University. FDA bioavailability submission document October 1981. 5. Lee DAH, et al. Gut 1980; 21: 215-218. Further information is available on request. **Stafford-Miller Ltd.**, Professional Relations Division, Hatfield, Herts. AL10 0NZ.

Antepsin[®]

Sucralfate

Mucoprotective ulcer healer



Non-systemic action

Fast pain relief

Excellent healing rates

Prolonged remission

Low incidence of side effects

Prescribing Information

Presentation Antepsin Tablets 1 gram are white, oblong, biconvex, uncoated tablets scored and embossed 1239 on one side and Ayerst on the other. Each tablet contains 1 gram sucralfate. **Uses** For the treatment of duodenal ulcer, gastric ulcer and chronic gastritis. **Dosage and Administration** For oral administration. **Adults** - Usual dose 1 gram 4 times a day. Maximum daily dose 8 grams. Four to six weeks treatment is usually needed for ulcer healing but up to twelve weeks may be necessary

in resistant cases. Antacids may be used as required for relief of pain. **Contra-indications, Precautions, Warnings, etc. Contra-Indications** There are no known contra-indications. **Precautions** 1. Concomitant administration with some oral anti-infectives such as tetracyclines may interfere with absorption of the latter. 2. The product should only be used with caution in patients with renal dysfunction. 3. As with all medicines, Antepsin should not be used in early pregnancy unless considered essential. **Side Effects** A low incidence of mild side effects, e.g. constipation, has been reported.

Legal Category POM. **Package Quantities** Antepsin 1 gram - Securitainers of 100. **Pharmaceutical Precautions** No special requirements for storage are necessary. **Product Licence Numbers** PL No. 0607/0045 PA No. 149/4/2. **Basic N.H.S. Price** Average daily cost 50p.



Ayerst Laboratories Ltd.
South Way, Andover, Hampshire SP10 5LT
Telephone: 0264 58711.
Distributors in Ireland: Ayerst Laboratories Ltd.,
765 South Circular Road, Islandbridge, Dublin 8.

* ANTEPSIN is a registered Trade Mark.

Further information is available on request to the Company.

Nature Plays a Dual Role in the Management of Irritable Bowel Syndrome

1. With the natural goodness of high-fibre foods

The irritable colon is affected by the amount of fibre in the diet. It is known that diets rich in high-fibre foods tend to normalize the function of the colon, which can result in softer, bulkier stools and a decrease in patient discomfort. Thus, a high-fibre diet is often considered basic therapy in the management of IBS.



2. With the natural efficacy of COLPERMIN

Colpermin provides natural relief to help the irritable bowel regain normal function. It has a powerful antispasmodic effect that relieves abdominal pain.

It is a naturally occurring carminative that relieves flatulence and gaseous distension. Enteric-coated capsules deliver relief direct to the site of action in the distal small bowel.

COLPERMIN™

(enteric-coated peppermint oil) CAPSULES



Henlow Trading Estate, Henlow, Beds. SG16 6DS

Presentation: Enteric-coated gelatine capsule. Each contains 0.2 ml standardised peppermint oil B.P. Ph. Eur. **Uses:** For the treatment of symptoms of discomfort and of abdominal colic and distension experienced by patients with irritable bowel syndrome. **Dosage and Administration:** One capsule three times a day, preferably before meals and taken with a small quantity of water. The capsules should not be taken immediately after food. The dose may be increased to two capsules, three times a day when discomfort is more severe. The capsules should be taken until symptoms

resolve, usually within one or two weeks. At times when symptoms are more persistent, the capsules can be continued for longer periods of between 2 to 3 months. There is no experience in the use of these capsules in children under the age of 15 years. **Contraindications, Warnings, etc.** **Precautions:** The capsule should not be broken or chewed. Patients who already suffer from heartburn, sometimes experience an exacerbation of these symptoms when taking the capsule. Treatment should be discontinued in these patients. **Adverse effects:** Heartburn, sensitivity

reactions to menthol which are rare, and include erythematous skin rash, headache, bradycardia, muscle tremor and ataxia. **Product Licence:** PL 0424/0009 **Basic NHS Cost:** £1058 per 100. UK and Foreign Patents pending. Colpermin is a trade mark of Tillotts Laboratories. Further information is available from Tillotts Laboratories, Henlow Trading Estate, Henlow, Beds.

European Patent No. 0015334. UK Patent No. 2006011



KEY MED

TOTAL RECALL

Is most of the data you record following an endoscopy consigned to oblivion? Or despatched with the patient? Or lost in another department? Using PEDRO as an integral part of your post-endoscopy routine will retain

full details for your own later use as well as providing hard copy for patients' notes and other interested parties. PEDRO'S total recall allows you to easily search for cases of interest, histological correlations and developing trends, and to improve management by recalling patients for follow-up, listing biopsy results and producing standard house-keeping audit figures. IT'S SAFE AND EASY TO USE.

- ★ No computer knowledge required – no keyboard codes.
- ★ Rapid data entry and short learning time (30 minutes).
- ★ Flexible and comprehensive – the data base is as simple or as detailed as you wish to make it.
- ★ Offers many features for analysis of clinical data and improvements in patient management.
- ★ Simplifies audit – justification for increased staffing and replacement of equipment.
- ★ Program design facilitates future expansion and enhancements.

Choosing a computer system is a long term investment – invest in the future with KeyMed – the company with a proven customer service record.

PATIENT ENDOSCOPY DATA & RECORDS ORGANISER

KEY MED

Specialised Services to Medicine
KeyMed House, Stock Road
Southend-on-Sea, Essex SS2 5QH,
England Tel (0702) 616333 (20 lines)
Telex: 995283 Telefax (0702) 65677

INTRODUCING Binary Cholelitholytic Therapy

For more effective dissolution and relief of symptoms of common bile duct gallstones,
use ROWACHOL in combination with chenodeoxycholic acid.¹

As the only adjuvant cholelitholytic agent containing monoterpenes derived from plant essential oils, ROWACHOL not only accelerates the dissolution of gallstones, but also permits reduction of the dose of chenodeoxycholic acid, thus reducing the potential for side effects.²

"...we reduced the chenodeoxycholic acid dose requirement by almost two-thirds;
this resulted in a great improvement in patient tolerance and reduced by half the total cost of treatment."²

ROWACHOL

(MENTHOL, PINENE, MENTHONE, CAMPHENE, BORNEOL, CINEOLE - COMPOUND OF CYCLIC MONOTERPENES) CAPSULES

ROWACHOL CAPSULES

PRESENTATION

Green enteric coated soft gelatin capsules, each containing:
Pinene 17mg, camphene 3mg, cineole 2mg, menthone 6mg, menthol 32mg,
borneol 5mg.

USES

Adjuvant therapy for the dispersal, dissolution and/or expulsion of stones in the common bile duct. To be used in combination with chenodeoxycholic acid for oral administration.

Adult dose: 1-2 capsules three times a day before meals. There is no dose recommendation for children.

CONTRAINDICATIONS, WARNINGS, ETC.

Caution should be used in patients receiving oral anti-coagulants or other agents metabolised by the liver, where the dose is critical.

Reduced cholesterol intake in the diet is advisable. Although no teratogenic effects have been reported, Rowachol should not be given in the first trimester of pregnancy.

BASIC NHS PRICE
50 - \$5.95

LICENCE HOLDER

Rowa Ltd, Bantry Co. Cork, Ireland
PL 0007 0002

ABBREVIATED PRESCRIBING INFORMATION

ROWACHOL LIQUID

PRESENTATION

Pale yellow liquid containing (in olive oil) v/v: menthol 32%, menthone 6%, pinene 17%, borneol 5%, cineole 2%, camphene 5%.

USES

Cholelithiasis, biliary and hepatic disorders.

DOSAGE AND ADMINISTRATION

For oral administration. Adult dose: 3-5 drops four or five times daily. No dose recommendation for children.

CONTRAINDICATIONS, WARNINGS, ETC.

Caution should be used in patients receiving oral anti-coagulants, or other agents metabolised by the liver, where the dose is critical.

Reduced cholesterol intake in the diet is advisable. Although no teratogenic effects have been reported, Rowachol should not be given in the first trimester of pregnancy.

Adverse effects: Eructation and a taste of peppermint can occasionally occur. Very occasionally, soreness of the mouth, or even buccal ulceration have been reported; these effects disappear on withdrawal of the drug.

BASIC NHS PRICE

10ml dropper bottle: \$5.70

LICENCE HOLDER

Rowa Ltd, Bantry Co. Cork, Ireland
PLR 0531/6286

REFERENCES:

1. Tills WR, et al. 'Oral dissolution therapy - a valid option in management of biliary duct stones'. *Gastroenterology*, in press.


2. Tills WR, Bell GD, Middleton B, et al. 'Adjunct to bile acid treatment for gallstone dissolution. Low dose chenodeoxycholic acid combined with a terpene preparation'. *BMJ* 1981; 282: 611-612.

Further information is available

on request from
Tillotts Laboratories,
Henlow Trading Estate,
Henlow Beds. SG16 6DS
Telephone:
0462 813933
Telex: 82313

Tillotts
LABORATORIES

The liquid life-line



for the intensive care patient



Administration of Fortison feeding is
most easily accomplished using
the Fortison Feeding Set (gastric drip line)
and the Fortison Tube.

For further information contact—
Cow & Gate Limited, Clinical Products Division,
Cow & Gate House, Trowbridge, Wiltshire, BA14 8YX.
Telephone: Trowbridge 02214 68381.



Renaissance

Mediaeval Crusades



Era of Richard III

Bodily defence still relies on shields


NOW! A natural mucosal shield helps heal peptic ulcers!

CAVED-S* does what no other ulcer therapy can do: it increases the number of mucus-secreting cells¹ with virtually no side effects.² This protects the gastric mucosal barrier against damaging agents^{3,4,5} and reduces ulcer recurrence.⁶

An 88% healing rate in 12 weeks⁷ has been reported. Studies also confirm that CAVED-S offers comparable efficacy to cimetidine in healing gastric ulcers⁷ and comparable efficacy to ranitidine in healing duodenal ulcers.⁶

REFERENCES:

1. Van Marle J, Aarsen PN, Lind A, et al: Deglycyrrhizinised liquorice (DGL) and the renewal of rat stomach epithelium. *Eur J Pharmacol* 72:219-225, 1981. 2. Cooke WM, Baron JH: Metabolic studies of deglycyrrhizinised liquorice in two patients with gastric ulcer. *Digestion* 4:264-268, 1971. 3. Rees WDW, Rhodes J, Wright JE, et al: Effect of deglycyrrhizinised liquorice on gastric mucosal damage by aspirin. *Scand J Gastroenterol* 14:605-607, 1979. 4. Morgan RJ, Nelson LM, Russell RI, et al: The effect of deglycyrrhizinised liquorice on the occurrence of aspirin and aspirin plus bile acid-induced gastric lesions, and aspirin absorption in rats, abstracted.



CAVED-S®

(deglycyrrhizinised liquorice, alum hydrox gel, mag carb, sod bic)

"The Mucosal Shield" for peptic ulcers



Henlow Trading Estate, Henlow, Bedfordshire. SG16 6DS.
Telephone 0462 813933 Telex: 82313 Tillab G.

PRESCRIBING INFORMATION

Presentation:

Brown tablets embossed 'CAVED-S', each containing:
Deglycyrrhizinised Liquorice 380 mg
Dried Aluminum hydroxide gel 100 mg
Magnesium carbonate 200 mg
Sodium bicarbonate 100 mg

Indications:

For the treatment of peptic ulcer and other allied conditions.

Dosage and Administration:

Adult dose for gastric ulcer:
2 tablets 3 times a day between meals.

Adult dose for duodenal ulcer:
Increase to 2 tablets 6 times a day between meals when necessary.

Prophylactic dose:

Gastric ulcer:

1 tablet 3 times a day, between meals.

Duodenal ulcer:

2 tablets 3 times a day, between meals.

Children's dosage 10-14 years:

half adult dose.

The tablets should be lightly chewed and swallowed with a drink of water, but in exceptional cases of objection to taste, the tablets should be broken into a few pieces and then swallowed with a drink of water. No additional antacids are necessary.

Contra-indications, warnings, etc:

Rare cases of mild diarrhoea can occur. No other side-effects have been reported.

Caved-S should be used with caution

in pregnancy.

Basic NHS Price:

60's—£2.83

240's—£10.12

600's—£22.76

PL0424/5000.



Gastroenterology 82:1134, 1982. 5. Morris TI, Calcraft BJ, Rhodes J, et al: Effect of a deglycyrrhizinised liquorice compound in the gastric mucosal barrier of the dog. *Digestion* 11:355-363, 1974. 6. McAdam WAF, Morgan AC, Pacsoo C, et al: A comparison between ranitidine and Caved-S in duodenal ulcer treatment, abstracted. Proceedings, World Congress of Gastroenterology, Stockholm, June 1982. 7. Morgan AC, McAdam WAF, Pacsoo C: Comparison between cimetidine and Caved-S in the treatment of gastric ulceration, and subsequent maintenance therapy. *Gut* 23:545-551, 1982.


SALAZOPYRIN[®] EN

sulphasalazine

HAS TOLERABILITY ALL WRAPPED UP

"Patients in whom sulfasalazine induces dyspeptic symptoms alone can be given EN Salazopyrin (entero-soluble) instead, and no more than 5% of these patients will be so troubled by dyspepsia that the treatment has to be discontinued."

Nielsen, O.H., Scand. J. Gastroenterol., 1982, 17: 389



Get them into the
SALAZOPYRIN habit
DAY AFTER DAY AFTER YEAR
500mg q.i.d. in ulcerative colitis

PRESCRIBING INFORMATION

Dosage and Administration Plain or EN Tabs. In acute/moderate attacks 2-4 tablets 4 times a day. If severe attacks give steroids also. Gradually reduce dose after 2-3 weeks to 3-4 tabs. day given indifferently. Suppositories: Two morning and night reducing dose after 3 weeks with improvement. Enema: One to be given at bedtime. Preparation contains adult dose.

Contra-indications Sensitivity to salicylates and sulfonamides. Infants under 2 years. Enema: Sensitivity to parabens.

Adverse Reactions Side effects common to salicylates or sulfonamides may occur. Most commonly these are nausea, loss of appetite and raised temperature which may be relieved on reduction of dose, use of EN tablets, enema or suppositories. If serious reactions occur the drug should be discontinued. Rare Adverse Reactions: Haematological: haemolytic anaemia, agranulocytosis, aplastic anaemia. Hypersensitivity: eg rash, fever. Gastrointestinal: eg stomatitis, impaired folate uptake. C.N.S.: eg peripheral neuropathy. Fertility: eg reversible oligospermia. Renal: eg proteinuria, crystalluria. Also: Stevens-Johnson syndrome and lung complications; eg fibrosing aveculitis.

Precautions Care in porphyria, allergic, renal or hepatic disease. Glucose 6-PD deficiency. Blood checks initially and periodically.

Pregnancy and Lactation While the ingestion of drugs in these situations may be undesirable, the severe exacerbations of the disease which can occur commends the continuance of therapy. Long clinical usage and experimental studies have failed to reveal teratogenic or fetotoxic hazards. The amounts of drug present in the milk should not present a risk to a healthy infant.

Package and Price Plain Tablets (0.5g) 100 & 500. £6.70 for 100. EN Tablets (0.5g) 100 & 500. £8.70 for 100. Suppositories (0.5g) 10 & 50. £2.80 for 10. Enemas (3.0g) 7, £12.10 for 7.


Product Licence Numbers Plain Tablets 0009/5006. EN Tablets 0009/5007. Suppositories 0009/5006. Enema 0009/5009.



Further information is available on request. Pharmacia Limited, Pharmacia House, Midsummer Boulevard, Milton Keynes MK9 3HP. Telephone Milton Keynes (0908) 681101.

LET GENTLE
SMOOTH

SPASMONAL

Alverine citrate 

SMOOTHES AWAY GUT SPASM.

Until now, no antispasmodic has been reliably assessed directly within the sigmoid colon . . .

SPASMONAL HAS. . . . and demonstrated to produce a significant reduction in colonic motor activity . . .²

SPASMONAL HAS. . . . not only reducing the spasm,² but also relieving pain and cramps . . .

SPASMONAL HAS. . . . and, because Spasmonal acts selectively, minimising side-effects, the only effect it has is the one you want for your patients.

Prescribing Information Presentation: Blue/grey opaque hard gelatin capsules each containing 60 mg Alverine Citrate USNF XIII. **Uses:** Selective smooth muscle spasmolytic. **Dosage and administration:** Adults 1 or 2 capsules one to three times daily, orally. No specific dosage recommendations can be made for children. **Contra-indications, warnings, etc.:** Nil. **Pharmaceutical precautions:** Store in a cool dry place. **Legal category:** P **Package quantities:** 100 capsules. **Basic NHS cost:** £6.98. **Further information:** Alverine citrate is a synthetic, non-narcotic, non-habit forming spasmolytic of a low order of toxicity in comparison with other anti-spasmodics. It is related to (but more than twice as active as) papaverine, and has a specific effect on the smooth muscle of the intestine and uterus, but not on those of the respiratory or cardiovascular system. **Product licence number:** 0322/5014. **References:** 1. Trotman, I.F. (awaiting publication). 2. Evangelista, I (1966) West. Med. 3, 49.



Norgine Limited,
116-120 London Road,
Oxford OX3 9BA

'Norgine' and 'Spasmonal' are trademarks.

318UK/17A/OSH/1U

ABC OF COMPUTING

A J ASBURY

Although computers are being widely used in medicine, their possibilities and limitations are still not clear to many potential users. This book, aimed at the non-expert, describes some of the uses of computers in medicine; because most doctors' involvement will be indirect, liaising with computer experts rather than designing systems themselves, the book concentrates on concepts rather than detailed descriptions of how computers work. It provides a useful introduction for the doctor who wants to know how computers can contribute to his practice of medicine.

Price: Inland £5.75; Overseas £8.00/USA \$14.00
(Inland £5.25; Overseas £7.50/USA \$13.00 to BMA members)
including postage by air overseas

Payment must be enclosed with order

Order your copy now

From: The Publisher, British Medical Journal
BMA House, Tavistock Square, London WC1H 9JR
or any leading bookseller

Zeitschrift für **GASTROENTEROLOGIE**

Organ der
Deutschen Gesellschaft für Verdauungs- und Stoffwechselkrankheiten,
der Deutschen Gesellschaft für gastroenterologische Endoskopie und
der Österreichischen Gesellschaft für Gastroenterologie

The highly specialized periodical
for gastro-enterologists and internists;

Top level original papers
relating to stomach, liver, pancreas and in-
testines;

Comprehensive information
on the advancements made in morphology,
endoscopy and roentgenology;

Extensive bibliography;

Conference papers.

Subscription one year (twelve issues) DM 114.-. postage to be added:



DEMETER VERLAG D-8032 GRÄFELFING
Tel.: (089) 85 20 33, Telex 05-24 068 delta d