


in the treatment of
peptic ulcer

ROTER

has particular
advantages





ROTER achieves a high cure rate and often gives effective control of even refractory chronic cases which have resisted other treatments.

ROTER in many cases obviates the need for hospitalization or surgical intervention.

ROTER helps patients to keep ambulant and symptom-free with a minimum of medication and with few, if any, dietary restrictions.

ROTER in general practice enables the physician to take adequate care of his gastric cases without frequent or prolonged supervision.

ROTER has no known side effects.

ROTER requires no supplementary medication such as sedatives or anti-spasmodics.

ROTER shows a saving in cost when compared with conventional methods of treatment.

REFERENCES

Therapie der Gegenwart, 1953, 3, 108
Gazz. Med. Ital., 1954, 113, 343
Die Medizinische, 1955, 4, 163
British Medical Journal, 1955, 2, 827
C.R.S. Biol., 1956, 150, 1019
Am J. Gastroenterology, 1957, 28, 439
Wiener Med. W., 1958, 108, 218
Ärztliche Praxis, 1958, 10, 766
Journal Lancet, 1960, 80, 37

FORMULA

Each Roter tablet contains:
Bismuth Subnitrate Roter (350 mg.), Magnesium
Carbonate (400 mg.), Sodium Bicarbonate
(200 mg.) and Cortex Rhamni Frangulae (25 mg.).

PACKINGS OF ROTER TABLETS

Tins of 40 and 120, also dispensing sizes,
360 and 720 (P.T. exempt). Basic N.H.S.
price: Under 1/- for 10 tablets.

ROTER TABLETS are in Category "S". Literature on Request

F.A.I.R. LABORATORIES LTD · TWICKENHAM · MIDDLESEX



People with worrying and exacting jobs are prone to gastric disorders.

For such patients 'Stelabid' is indicated. 'Tyrimide', the powerful and long-acting anticholinergic component in 'Stelabid', relieves pain in gastro-intestinal disorders; it inhibits hypermotility and spasm; it reduces excessive gastric secretion.

The 'Stelazine' constituent in 'Stelabid' prevents nausea and vomiting; it also allays the nervous stress and strain which can cause and aggravate gastric disturbances.

Exceptionally long-acting, 'Stelabid' ensures night and day relief of pain on a convenient b.d. dosage regimen.

Each tablet of 'Stelabid' contains 1 mg. 'Stelazine' (brand of trifluoperazine) and 5 mg. 'Tyrimide' (brand of isopropamide)

STELABID

— relieves physical pain and nervous strain in peptic ulcer and other gastrointestinal disorders

**SK
&F**


**Smith Kline & French
Laboratories Ltd
Welwyn Garden City, Herts**

'Stelabid', 'Stelazine' and 'Tyrimide' are trade marks
Brit. Pat. Nos. 772921, 813861, 857547
SBD:PA12

a more convenient and economical
presentation of hydrocortisone
for ulcerative colitis

EFCORTELAN SOLUTION

TRADE MARK




Given as a nightly rectal drip, soluble hydrocortisone (Efcortelan) has been demonstrated to afford rapid alleviation of symptoms of ulcerative colitis in most patients. Symptomatic relief is accompanied by an improvement in the sigmoidoscopic findings and in the colonic mucosa.

The method of treatment is quite suitable for the patient to carry out himself provided the attack is not severe.

With soluble Efcortelan, hormonal side-effects are avoided. If further attacks occur they can usually be

reduced by another course of treatment.



Only one of the new Efcortelan Solution Tablets in the appropriate amount of normal saline (100-120 ml) is needed to prepare the retention enema. It is therefore the ideal presentation for home treatment.

EFCORTELAN SOLUTION TABLETS

Foil strip of 7 tablets (one week's treatment). Each tablet contains 100 mg hydrocortisone (as water soluble sodium succinate).

Subsidiary Companies or Agents in most countries

**GLAXO LABORATORIES LTD.
GREENFORD, MIDDLESEX
BYRon 3434**

Uniquely effective in
antibacterial
and antiamoebic
therapy

Humatin **KAPSEALS**

HUMATIN* is unique among broad-spectrum antibiotics in that it is highly effective against intestinal amoebiasis as well as many types of bacterial enteric infections. HUMATIN (Paromomycin, P.D. & CO.) has also been found useful in the management of hepatic coma and in suppressing, pre-operatively, the normal intestinal microflora.

Available in Kapseals* each containing the sulphate equivalent to 250 mg. paromomycin base. In bottles of 16 and 100.

Now also available as Humatin Syrup in bottles of 60 ml. and 500 ml.; each 5 ml. containing paromomycin sulphate equivalent to 125 mg. paromomycin base.

*TRADE MARK

PARKE-DAVIS

WPS 1102



PARKE, DAVIS, HOUNSLOW, MIDDLESEX. Tel: HOUnslow 2361

ASPIRIN IS SAFER WHEN IT IS SOLUBLE

*Soluble aspirin is safer
when it is stable*



N.H.S. Basic Price 12/6 for 500 tablets in foils.

SOLPRIN provides soluble aspirin in its most stable form and dissolves rapidly in water to form a solution of calcium acetylsalicylate.

SOLPRIN is easy to take and palatable. It has proved rapid and reliable in its analgesic effect and is extremely well tolerated, even in prolonged and massive dosage.

SOLPRIN is not advertised to the public and is available only on prescription. Clinical samples and literature will be supplied on request.

SOLPRIN

REGD.

Soluble and substantially neutral

RECKITT & SONS LIMITED, PHARMACEUTICAL DEPARTMENT, HULL.

FOR
CONFIDENT DIAGNOSIS
IN GASTROENTEROLOGY

SECRETIN and PANCREOZYMIN

These two preparations can play important roles as diagnostic agents in gastroenterology. They may be used with confidence in the serum enzyme test and the duodenal aspiration test.

SECRETIN

is supplied in 10 cc rubber-capped vials each containing 80-100 units. The potency is declared on each vial following biological assay.

PANCREOZYMIN


is supplied in 20 cc rubber-capped vials each containing 100 units.

Please write for further information and supplies to:

BOOTS PURE DRUG COMPANY LIMITED
STATION STREET · NOTTINGHAM · ENGLAND



**A BOOTS
MEDICAL
PRODUCT**



**NOW
IN**




**JUST
TEAR
OPEN!**

*

SURGICEL

ABSORBABLE
HAEMOSTAT



Oozing from the gall bladder bed can be readily controlled by SURGICEL* which conforms closely to the exposed contour and intimately contacts the bleeding surface.

A MAJOR ADVANCE IN HAEMOSTASIS

effective control of difficult bleeding

SURGICEL haemostatic gauze completely controls capillary and venous oozing. Its use increases the range and safety of surgical procedures on highly vascular areas.

easily handled

SURGICEL is more easily handled than other agents in this field. It does not fragment, can be sutured in position and adheres closely to the site of bleeding but does not cling to gloves or instruments.

completely absorbed


SURGICEL is made from oxidized regenerated cellulose. It is absorbed with minimum tissue response and may be safely left *in situ*. No case of intolerance or sensitivity has been recorded.

Johnson & Johnson PRODUCTS TO MATCH THE SURGEON'S SKILL
(Gt Britain) Ltd

HOSPITAL DIVISION · SLOUGH · BUCKS

*a trade mark of Johnson & Johnson JJ 71

**GASTRIC ACID
SECRETION
BACK TO NORMAL
WITH**



NACTON

REGD. TRADE MARK

**NACTON GIVES RELIEF FROM PAIN IN
PEPTIC ULCER AND HYPERACIDITY**


- * 50%—75% reduction of gastric acid secretion
- * Activity prolonged for as much as 8 hours
- * Selective action
- * No disturbing side effects

Each Nacton tablet contains 2 mg. of (1-methyl-2-pyrrolidyl) methyl benzilate methylmethosulphate.



NACTON is a product of British Research
at **BEECHAM RESEARCH Laboratories Ltd.**
Brentford, England

**ALL YOUR
EGGS . . .**



IN ONE BOTTLE

Not really, because Aminosol is a casein hydrolysate. It does, however, provide adequate protein parenterally where oral nutrition

is impracticable.

Supplied as Aminosol Glucose Vitrum (*Aminosol G.V.*)

307 calories per litre and

Aminosol Fructose Ethanol Vitrum (*Aminosol G.E.V.*)

875 calories per litre

In 500 and 1000 ml. bottles. Aminosol Glucose Vitrum also provided in 100 ml. paediatric size. Further information on request

AMINOSOL

Vitrum

PAINES & BYRNE LTD., PABYRN LABORATORIES, GREENFORD, MIDDX.

to aid the
management of
peptic ulcer

PORTYN* produces marked depression of gastro-intestinal spasm, gastric secretion and hypermotility, and, at effective dosage levels, the incidence of side-effects is unusually low.

PORTYN is thus recommended as an excellent adjunct in the symptomatic control of peptic ulcer and gastro-intestinal spasm.

PORTYN

potent anticholinergic agent

PORTYN Kapseals each contain 10 mg. benzilonium bromide. Initial adult dose -- 1 Kapseal three times daily.

* TRADE MARK

PARKE-DAVIS

HOUNSLOW, MIDDLESEX. TELEPHONE: HOUNslow 2361

Of all the array of procedures available for the treatment of peptic ulceration mention must be made of the eminence of propantheline bromide.*

reduces gastric secretion
significantly

relieves pain rapidly

John Harrison

Pro-Banthine

standard anticholinergic in peptic ulcer therapy

AVAILABLE in 15 mg. tablets of propantheline bromide in packs of 100 and 1000.
also available: Pro-Banthine with Dartalan;
Pro-Banthine with Phenobarbitone, B. P.
BASIC N.H.S. COST 23s. 2d. and 217s. 10d. respectively.

**Brit. J. clin. Pract.* 11 452 (1957)


calms hypermotility quickly



G. D. Searle & Co. Ltd.

High Wycombe Bucks

transid
regd.
 polyaminostyrene



**for acid
 transfer**

the polymer for acid transfer

'Transid' is active only against excess acid; it also reduces pepsin activity. Otherwise, 'Transid' is completely inert; it does not interfere with digestion or absorption and cannot cause alkaloses, constipation or diarrhoea. Clinical experience suggests that 'Transid' may succeed in cases of peptic ulcer in which medical treatment with antacids has failed.

INDICATIONS:

Peptic ulcer; gastric hyperacidity from whatever cause; dyspepsia of pregnancy.

PACK AND BASIC NHS PRICE:

100 foil-pack tablets 10/—.


May we send you a copy of our Standard Publication on 'Transid'? It contains much more information than can be included in a journal advertisement.

DUNCAN FLOCKHART OF EDINBURGH

The Doctors' House

Duncan, Flockhart & Co Ltd Wheatfield Rd Edinburgh 11

ULCERATIVE COLITIS



In Ulcerative Colitis
extensive clinical
experience confirms

SALAZOPYRIN

BRAND OF SULPHASALAZINE
(SALICYLAZOSULPHAPYRIDINE)

to be a drug of
significance in to-day's
management of this
distressing disease.

The present investigation suggests that
salazopyrin is a useful drug. *Gut* 1:217 (1962)

Outstanding characteristics of Salazopyrin treatment are:

- Prompt remission of symptoms
- Bleeding and frequency of stools sharply reduced
- In many cases rectal mucosa heals within a month
- Abdominal cramp controlled
- Fever subsides, appetite improves
- Can be given safely over long periods of time

SALAZOPYRIN is also a safe and effective product in other forms of colitis and regional enteritis.

 PHARMACIA
UPPSALA SWEDEN

Distributed by Sandoz Chemicals Ltd., 90 Cl. W. Bedford Street, London, W.1.

When
infection
is at
hand



'Streptotriad'

trade mark

brand

TRI-SULPHONAMIDE/STREPTOMYCIN

against bacillary dysentery

for the treatment of the acute phase
and during the convalescent or chronic stage to prevent
the patient becoming a carrier.

Each 'Streptotriad' tablet and 3.6 ml. teaspoonful of prepared suspension contains:

Streptomycin (as sulphate B.P.) 65 mg.

Sulphamerazine B.P. 65 mg.

Sulphadiazine B.P. 100 mg.

Sulphathiazole B.P.C. 100 mg.



An M&B brand Medical Product

MANUFACTURED BY **MAY & BAKER LTD**

MA 8256

DISTRIBUTORS PHARMACEUTICAL SPECIALITIES (MAY & BAKER) LTD · DAGENHAM

THE BRITISH MEDICAL ASSOCIATION AND THE PROMOTION OF KNOWLEDGE

ONE of the foremost aims of the British Medical Association is the promotion of medical and allied sciences and each year a considerable portion of its income is spent in the attainment of this object through the publication of a wide range of general and specialist journals

The chief of the publications is the BRITISH MEDICAL JOURNAL with a weekly circulation of over 84,000 copies. The Journal serves as a comprehensive record of current developments and is a forum for discussion in every field of medical science and professional organization in this country

The List of SPECIALIST JOURNALS includes

ABSTRACTS OF WORLD MEDICINE *Published Monthly Annual Subscription £4 4s.*

BRITISH JOURNAL OF OPHTHALMOLOGY *Published Monthly Annual Subscription £6*

OPHTHALMIC LITERATURE *Published Quarterly Annual Subscription £4 4s.*

(Combined Annual Subscription for the two Ophthalmic Journals £9)

ARCHIVES OF DISEASE IN CHILDHOOD *Published Bi-monthly Annual Subscription £4 4s.*

BRITISH HEART JOURNAL *Published Bi-monthly Annual Subscription £4 4s.*

BRITISH JOURNAL OF PHARMACOLOGY AND CHEMOTHERAPY *Published Bi-monthly Annual Subscription £6 6s.*

JOURNAL OF CLINICAL PATHOLOGY *Published Bi-monthly Annual Subscription £4 4s.*

QUARTERLY JOURNALS *Uniform Annual Subscription £3*

ANNALS OF THE RHEUMATIC DISEASES

BRITISH JOURNAL OF INDUSTRIAL MEDICINE

BRITISH JOURNAL OF PREVENTIVE AND SOCIAL MEDICINE

BRITISH JOURNAL OF VENEREAL DISEASES

GUT

JOURNAL OF NEUROLOGY, NEUROSURGERY AND PSYCHIATRY

MEDICAL AND BIOLOGICAL ILLUSTRATION

THORAX

ANY QUESTIONS? (Series 3) Price 7s. 6d. (by post 8s. 3d.)

A selection of Questions and Answers from this popular weekly feature in the British Medical Journal

REFRESHER COURSE FOR GENERAL PRACTITIONERS *Second and Third Volumes Price 25s. each (by post—inland and overseas 26s. 9d.)*

The third volume of Refresher Course articles that have been printed in the British Medical Journal has now been published in book form. There are sixty chapters and each one has been revised and brought up to date by the author. A cumulative classified index of all three books in this series has been added

For further details please apply to

**The Publishing Manager, BRITISH MEDICAL ASSOCIATION
TAVISTOCK SQUARE, LONDON, W.C.1**

STAPHYLOCOCCAL SUPERINFECTIONS

"Patients in hospital who have had an operation, and a course of 'broad-spectrum' antibiotics, sometimes develop diarrhoea which may be severe and even fatal."

Lancet Annotations 25th February, 1961

ENPAC

LACTOBACILLUS ACIDOPHILUS PREPARATION

For the prophylaxis and treatment of staphylococcal super infections.

"It was found that the staphylococci increased in both groups at the beginning of antibiotic treatment. Staphylococci continued to increase in the patients taking the antibiotic alone, but in those taking 'ENPAC' there was a pronounced drop in numbers."

Gordon D., Macrae J., & Wheeler D. M. W. Lancet—May 4th, 1957. pp. 899-901

Samples and literature will be sent to the medical profession on request to:—

WILTS UNITED DAIRIES LIMITED · NUTRITION DEPARTMENT · TROWBRIDGE · WILTS

"This book will take its place as the reference volume on the surgery of the anus, rectum, and colon, both at home and overseas"—*British Journal of Surgery*.

Surgery of the Anus, Rectum and Colon

J. C. GOLIGHER, Ch.M., F.R.C.S.

Professor J. C. Goligher holds the chair of Surgery at the University of Leeds, and is Surgeon, General Infirmary at Leeds; formerly Surgeon St. Mary's Hospital and St. Mark's Hospital for Diseases of the Rectum and Colon, London.

This book, written especially for Surgeons in training and for young Consultants, comprehensively surveys contemporary knowledge of diseases of the rectum and colon, and at the same time provides a clear and detailed practical guide to the surgical management of these conditions. A special feature is the two chapters on developmental abnormalities of the anus and rectum and on Hirschsprung's Disease and related abnormalities by Mr. Harold Nixon, F.R.C.S., of the Hospital for Sick Children, Great Ormond Street, London.
896 pp. Crown 4to 5 colour plates 276 half-tone illustrations. £8 8s. net.


CASSELL MEDICAL BOOKS

'ALUHYDE'

for prompt, safe symptomatic medication in
hyperchlorhydria and peptic ulcer

'ALUHYDE' tablets present a new combination with these advantages...


'ALUHYDE' effects safe, reliable, non-systemic neutralization by means of two of the most preferred modern antacid adsorbents...



'ALUHYDE' permits of quick mental sedative-effect by its inclusion of a barbiturate recognized for minimal cumulative potential...



'ALUHYDE' provides antispasmodic action through its content of an agent long known for its favourable influence against hypermotility...



Formula—

Each tablet: Dried Aluminium Hydroxide Gel. B.P.C.	-	3.75 gr.
Magnesium trisilicate B.P.	-	3.75 gr.
Quinalbarbitone sodium B.P.	-	0.50 gr.
Tinct. Belladonnæ B.P.	-	3 m.

Packings—Regular packs of 50 ; bottles of 500 (dispensing)

*Samples available on personally signed request
of qualified physicians only*

MULTIPAX CHEMICALS LTD

121 NORWOOD ROAD, LONDON, S.E.24

Phone : TULse Hill 3276



IAS3