

Leading article

Helicobacter pylori and non-steroidal anti-inflammatory drugs: uncomfortable partners in peptic ulcer disease

Recent studies have suggested that *Helicobacter pylori* and non-steroidal anti-inflammatory drugs (NSAIDs) are capable of interfering with various protective mechanisms in the gastroduodenal mucosa. Whereas NSAIDs are recognised for their acid stimulating activity, the effect of *H pylori* on gastric acid secretion remains highly speculative despite its association with hypergastrinaemia. Both *H pylori* and NSAIDs have, however, been shown to influence the production rate and the quality of gastric cyclic AMP, the mucus layer, mucosal prostaglandins, blood flow, and platelet activating factor. Characteristic histological abnormalities have also been identified. Also NSAID related peptic ulcers seem to develop more commonly in patients infected with *H pylori* despite the apparent reduction in the prevalence of these organisms in chronic NSAID users, hence the uncomfortable relation between the two aetiological factors. It remains to be seen whether the eradication of *H pylori* would reduce the frequency of peptic ulcers induced by NSAIDs or prevent their recurrence.

Helicobacter pylori and NSAIDs are probably the commonest known exogenous factors in the aetiology of peptic ulcer disease. This paper describes the pathogenetic mechanisms common to *H pylori* and NSAIDs and discusses the possibility of a synergistic relation between them. The importance of such a relation, if present, relates to its potential to provide a new therapeutic approach to the common, but yet unresolved, problem of ulcers induced by NSAIDs.

Effect of NSAIDs and *H pylori* on gastric acid secretion

Because of the widely accepted importance of gastric acid in the pathogenesis of ulcers in general, it was only natural for many workers to investigate the possible effects of NSAIDs and, more recently, *H pylori* on acid secretion. NSAIDs (indomethacin and aspirin in particular) were found to increase basal and maximally stimulated gastric acid secretion^{1,3}; they seem to bypass the H₂ and muscarinic receptors and interact with secretagogues at a locus between the catalytic subunit of adenylate cyclase activation and the proton pump.⁴ The potentiation of secretagogue stimulated acid secretion by non-salicylate NSAIDs has also been found to be dependent on calcium.⁴

The situation is not so well defined in the case of *H pylori*. Because of its association with hypergastrinaemia, it was speculated that *H pylori* could increase the parietal cell mass that is characteristic of patients with duodenal ulcers.⁵ To date, the evidence for an increase in gastric acid secretion by *H pylori* has, however, been lacking.^{5,6} On the contrary, there is a consensus that acute exposure to *H pylori* causes hypochlorhydria.⁷⁻¹¹

Recent evidence indicates that hypergastrinaemia associated with *H pylori* might not be directly related to the function of the parietal cells,¹² the number of antral G cells,¹³ or to the bacterium's urease activity.¹⁴ It is more likely to be related to local inflammation¹³⁻¹⁵ or products of the T lymphocyte such as interleukin-2 and γ -interferon.¹⁶ Gastric

acid secretion in chronic NSAID users who are infected with *H pylori* has not been studied, but indomethacin has been shown to potentiate the inhibitory effect of *H pylori* protein on gastric fundic cyclic AMP, which in turn mediates acid secretion in vitro.¹⁷ This might explain, at least in part, the tendency of many arthritic patients to develop a degree of hypochlorhydria,¹⁸ and the limitations of acid inhibition in the management of NSAID related ulcers.^{19,20}

The mucus layer

The ability to change the characteristics of the gastric mucus layer is common to both NSAIDs and *H pylori*. Aspirin²¹ and indomethacin²² were found to inhibit mucus secretion. Aspirin can also increase pepsin mediated proteolysis of mucus, decrease mucus viscosity, and increase the permeability of mucus to hydrogen ion.²³ Indomethacin was shown to inhibit active bicarbonate secretion by the gastric mucosa.²⁴ It was also suggested that NSAIDs could cause disruption of the gastric mucosal barrier, which in turn allows back diffusion of hydrogen ion with its damaging results.²⁵⁻²⁷

Similarly, it has recently been shown that incubation of a culture filtrate of *H pylori* with gastric mucus could lead to a gradual loss of viscosity of mucus,²⁸ which might impair the ability of the mucus to retard the diffusion of hydrogen ions. It was concluded that the degenerative changes produced in the gastric mucus gel by *H pylori* might be a contributing factor to the pathogenesis of gastritis and peptic ulcers.²⁸

Mucosal prostaglandins

It is generally accepted that NSAIDs, with few exceptions,²⁹ are capable of suppressing gastric and duodenal mucosal prostaglandin (PG) synthesis, although no correlation has been found between the degree of inhibition of PG and the endoscopic abnormalities.^{29,30}

Because of its strong association with gastritis and neutrophilic infiltration, *H pylori* is expected, at least in theory, to stimulate PG production, as human neutrophils and macrophages are capable of synthesising PGs.³¹ Also gastritis – like any other inflammation – is associated with raised PG values irrespective of *H pylori* status.^{32,33} It was surprising, therefore, to find that patients with *H pylori* related gastritis had PG values similar to those without *H pylori*.³³⁻³⁵ This led to speculation that *H pylori* infection might result in at least a partial block in PG synthesis at the level of the neutrophils, mucosal cells, or both.³³ Indeed, some workers reported reduced concentrations of stable metabolites of PGI₂ in patients infected with *H pylori*,³⁶ but this was not confirmed by others.³³⁻³⁵ It was also interesting to find that, in vitro, the combination of indomethacin and *H pylori* culture filtrate reduced gastric antral mucosal PGE₂ and epithelial viability to a greater degree than indomethacin alone.³⁷ This in turn might suggest a synergistic relation between *H pylori* and NSAIDs in causing mucosal damage.

Mucosal blood flow

Several studies have shown that short courses of NSAIDs could reduce gastric mucosal blood flow in laboratory animals.^{2 38 39} The situation is not clear in patients on long term NSAIDs, who could have other factors that might alter blood flow. Also, little is known about the duodenal micro-circulation in such patients. It has been shown, however, that duodenal, but not gastric, mucosal blood flow is lowest in NSAID patients who smoke, or those with duodenal ulcers or *H pylori*⁴⁰; no correlation could be found with age or sex. Further analysis of the data in this study⁴⁰ showed that the suppressive effect of smoking and *H pylori* on duodenal blood flow in NSAID patients was independent of ulceration and other demographic variables. Although such findings do not dispute the capacity of NSAIDs to lower mucosal blood flow in general, they might represent another aspect of the synergistic relation thought to exist between NSAIDs and *H pylori*, at least for duodenal blood flow.

Histological and endoscopic abnormalities

Active chronic gastritis is the classic histological picture found in association with *H pylori* in most infected subjects. It is best remembered as the main abnormality seen after the deliberate ingestion of the organisms to fulfil Koch's postulates for the pathogenicity of *H pylori*,^{7 8} and in epidemic gastritis.⁹ It is also the commonest histological finding in patients treated with NSAIDs.⁴¹ Neutrophilic infiltration, common to most cases of NSAID and *H pylori* gastritis, could be related to the capacity of NSAIDs and *H pylori* to stimulate the production of platelet activating factor.^{42 43} As well as its ulcerogenic actions,⁴² platelet activating factor might be involved in the aggregation and activation of neutrophils,⁴⁴ which might subsequently contribute to ulcer formation: neutropenic rats are less likely to develop gastric mucosal injury from indomethacin or naproxen.^{44 45} These interesting findings might not apply, however, to patients with chemical gastritis, another histological abnormality characteristic of chronic NSAID intake.^{41 46} Not unlike bile reflux gastritis in patients with a history of gastroduodenal surgery,⁴⁷ it is characterised by foveolar hyperplasia, vasodilatation, oedema, lack of inflammatory cells, and the presence of muscle fibres in the lamina propria.^{41 46 49} *H pylori* is very rare in cases of chemical gastritis,^{41 46 47} and the histological picture can be found in about 25% of chronic NSAID users.^{41 46} The importance of such an entity stems from the positive correlation between the histological chemical scores and the degree of endoscopic damage: peptic ulcers related to NSAIDs and measuring 5 mm in diameter were found more commonly in patients with chemical gastritis or gastritis associated with *H pylori* than in those without either of these conditions.⁴¹ This is in agreement with other studies,^{50 51} that found a greater number of gastric ulcers in NSAID patients infected with *H pylori* compared with patients either taking NSAIDs or infected with *H pylori*. The prevalence of NSAID related ulcers in *H pylori* positive patients has also been found to be twice that in *H pylori* negative subjects, but the differences were not significant due to the small numbers of patients and ulcers studied.^{52 53} Graham *et al*⁵² found a greater number of submucosal haemorrhages and erosions in *H pylori* negative NSAID users, unlike the results of another report, which suggested that the presence of *H pylori* had no influence on the prevalence of such lesions after the acute administration of naproxen or aspirin in healthy volunteers.⁵⁴ Also, Shallcross *et al*⁵³ found more duodenal than gastric ulcers in their NSAID patients, which is not typical of NSAID related damage, and could be explained by the fact that they included in their study patients who were able to stop their NSAID within a month before endoscopy. This would in

turn raise concern about the suitability of such subjects for analysis and their need for and compliance with NSAID intake, which can determine the extent of peptic damage. A more recent study in asymptomatic NSAID users from the community has found that *H pylori* infection was not related to the severity of the endoscopic findings, but to higher degrees of gastritis,⁵⁵ which in the long term might progress to ulceration.⁵⁶

The interpretation of the findings related to submucosal haemorrhages or erosions is made difficult by the chronic nature of *H pylori* infection, its association with ulcers, and the lack of evidence that such minor lesions can really progress to ulcers. Another possible explanation for the differences in the findings of these studies^{41 50 55} is the failure to correct for the presence of chemical gastritis, which appears to act as an independent factor, and the use of serological tests of unproved value in patients treated with NSAIDs. Diagnostic titres of *H pylori* IgG antibodies could still be detected in patients with chemical gastritis, with its high prevalence of NSAID ulcers, despite the failure to confirm the presence of the organisms by culture, histology, or urease activity.^{57 58} Also serology might not be reliable enough in predicting the presence of *H pylori* or ulcers in NSAID patients, unlike other categories of patients,^{59 60} because: firstly, NSAID ulcers can be completely without symptoms^{61 62}; secondly, the specificity of some IgG tests might be low in patients receiving NSAIDs^{57 58}; thirdly, despite the apparent synergistic relation between *H pylori* and NSAIDs, the intake of such drugs might be associated with a reduced prevalence of *H pylori* infection^{41 57 63 65}; this could be due to a direct toxic action, or indirectly related to the increase in acid secretion^{1 4} and the interference with the mucus layer^{21 27} rendering the natural habitat of *H pylori* organisms unsuitable for their survival.

The virulence of *H pylori* strains might also explain some of the differences in the reported rates of ulcer formation. Western blotting analysis of systemic IgG or IgA responses to *H pylori* has shown the antigenicity of 110-120 kDa, 89 kDa, 61 kDa, 54 kDa, and 31 kDa proteins, although there is substantial variability among subjects.^{66 70} The 120 kDa protein, which is recognised systemically in 83% of *H pylori* positive subjects, is a surface protein not expressed in some *H pylori* strains.⁶⁸

It has been suggested that 120 kDa positive strains have pathogenetic features associated with active gastritis and peptic ulcerations.⁷⁰ Infection with 120 kDa negative strains might explain why peptic ulceration develops in only a proportion of subjects infected with *H pylori*.⁷⁰ A vacuating cytopathic agent has also been described in some strains of *H pylori*.⁷¹ It is not, however, known whether these important findings would apply to patients receiving long term NSAIDs.

Therapeutic considerations

The role of eradicating *H pylori* in the healing or the prevention of NSAID related peptic ulcers has not been studied. As well as their antibacterial activity, however, agents currently used in the treatment of *H pylori* infection were previously found to possess properties relevant to the mechanisms of NSAID induced mucosal damage. Colloidal bismuth subcitrate was shown to stimulate gastric and duodenal alkaline secretion through a prostaglandin dependent mechanism.⁷² The same agent was also found to have a protective action against aspirin induced micro-bleeding, and this protection occurred despite a pronounced suppression of mucosal prostaglandin production.⁷³ Also, it was shown that the ulcerogenic effect of NSAIDs could be reduced in animals by treatment with antibiotics,⁷⁴ and that germ free animals were resistant to indomethacin induced intestinal lesion.^{75 76}

The common nature of NSAID related ulcers in association with *H pylori* colonisation, especially in elderly patients, should act as an incentive to devise an effective therapeutic approach. In 1983, enough NSAIDs were made available to treat almost three million people daily in the United States,⁷⁷ and 22 million prescriptions were issued for such agents in the United Kingdom.⁷⁸

About half of the NSAIDs prescribed were for persons over 60 years old.⁷⁸ *H pylori* infection can be identified in about 80% of people of this age group⁷⁹ although in the presence of NSAIDs the prevalence of *H pylori* might fall to 30% to 50%, according to data collected at our unit between 1987 and 1992.

It is estimated, therefore, that between 25% and 40% of all NSAID users and 30–50% of elderly patients taking NSAIDs could be infected with *H pylori* and, as a result, are at special risk of developing peptic ulcers. The management of ulcers in this large number of patients is not only essential but might also need to be different from that of other types of ulcers. Its effect on ulcer complications, especially in elderly patients, also needs to be clarified.

Despite the evidence for a low prevalence of *H pylori* in chronic NSAID users, it has to be emphasised that the possible use of NSAIDs to eradicate *H pylori* should not be considered, knowing that such eradication might involve the impairment of the mucus layer with its damaging consequences.

In conclusion, the evidence for a synergistic relation between NSAIDs and *H pylori*, despite being contested, should not be ignored because of its potential therapeutic implications. Also, the suggestion that NSAIDs might influence the prevalence of *H pylori* would indicate that the two factors are uncomfortable partners in peptic ulcer disease.

A S TAHA
R I RUSSELL

Gastroenterology Unit,
Royal Infirmary,
Glasgow G31 2ER

- Feldman M, Coltrac TJ. Effect of indomethacin on gastric acid and bicarbonate secretion in humans. *Gastroenterology* 1984; **87**: 1339–43.
- Gerkens JR, Shand DG, Flexner C, Nies A, Oates J, Data J. Effect of indomethacin and aspirin on gastric blood flow and acid secretion. *J Pharmacol Exp Ther* 1977; **203**: 646–52.
- Levine RA, Schwartzel EH. Effect of indomethacin on basal and histamine stimulated human gastric acid secretion. *Gut* 1984; **25**: 718–22.
- Levine RA, Jyotirmoy N, King RL. Non-salicylate non-steroidal anti-inflammatory drugs augment prestimulated acid secretion in rabbit parietal cells. Investigations of the mechanisms of action. *Gastroenterology* 1991; **101**: 756–65.
- Smith JTL, Pounder RE, Nwokolo CU, Lanzon-Miller S, Evans DG, Graham DY, Evans (Jr) DJ. Inappropriate hypergastrinaemia in asymptomatic healthy subjects infected with *Helicobacter pylori*. *Gut* 1990; **31**: 522–25.
- Wagner S, Gladziwa U, Haruma K, Verrenttrapp M, Gebel M. Effect of *Helicobacter pylori* infection on 24 hour intragastric acidity in patients with gastritis and duodenal ulcer. *Gut* 1992; **33**: 1024–8.
- Marshall BJ, Armstrong JA, McGeachie DB, Glancy RJ. Attempt to fulfil Koch's postulates for pyloric *Campylobacter*. *Med J Aust* 1985; **142**: 436–7.
- Morris A, Nicholson G. Ingestion of *Campylobacter pyloridis* causes gastritis and raised fasting gastric pH. *Am J Gastroenterol* 1987; **82**: 192–9.
- Peterson W, Lee E, Skoghund M. The role of *Campylobacter pyloridis* in epidemic gastritis with hypochlorhydria. *Gastroenterology* 1987; **92**: A1575.
- Graham DY, Smith JL, Alpert LC, Yoshimura HH. Epidemic achlorhydria is not viral but is caused by *Campylobacter pyloridis*. *Gastroenterology* 1987; **92**: A1412.
- Cave DR, Vargas M. Effect of *Campylobacter pylori* protein on acid secretion by parietal cells. *Lancet* 1989; **ii**: 187–9.
- Chittajallu RS, Harwood J, Dorrian CA, McColl KEL. Is *Helicobacter pylori* related hypergastrinaemia due to the bacterium inhibiting parietal cell function. *Gut* 1991; **32**: A1206.
- Graham DY, Lew GM, Lechago J. *Helicobacter pylori* related increase in plasma gastrin: What is the status of the antral G cells in man? *Gastroenterology* 1991; **100**: A75.
- Chittajallu RS, Dorrian CA, Neithercut WD, Dahill S, McColl KEL. Is *Helicobacter pylori* associated hypergastrinaemia due to the bacterium's urease activity or the antral gastritis? *Gut* 1991; **32**: 1286–90.
- Wyatt JI, Rathbone BJ, Green DM, Primrose J. Raised fasting serum gastrin in chronic gastritis is independent of *Campylobacter pylori* status and duodenal ulceration. *Gut* 1989; **30**: A1483.
- Teichmann RK, Pratschke E, Grab J, Hammer C, Brendel W. Gastrin release by interleukin-2 and gamma-interferon in vitro. *Can J Physiol Pharmacol* 1986; **64** (suppl): 62.
- Taha AS, Fraser WD, Kelly RW, Gemmell CG, Lee FD, Russell RI. Inhibition of human gastric cyclic AMP production by *Helicobacter pylori* protein – possible involvement of mucosal prostaglandin E₂. *Aliment Pharmacol Therap* 1991; **5**: 379–89.
- De Witte RJ, Geerdink PJ, Lamers CB, Boerbooms AM, Van der Korst JK. Hypochlorhydria and hypergastrinaemia in rheumatoid arthritis. *Ann Rheum Dis* 1979; **38**: 14–7.
- Ehsannullah RSB, Page MC, Tildesley G, Wood JR. Prevention of gastroduodenal damage induced by non-steroidal anti-inflammatory drugs: controlled trial of ranitidine *BMJ* 1988; **297**: 1017–21.
- Lanza FL, Aspinall RL, Swabb EA, Davies RE, Rack MF, Rubin A. Double-blind, placebo-controlled endoscopic comparison of the mucosal protective effects of misoprostol versus cimetidine on tolmetin-induced mucosal injury to the stomach and duodenum. *Gastroenterology* 1988; **95**: 284–94.
- Menguy R, Masters YF. Effect of aspirin on gastric mucus secretion. *Surg Gynaecol Obstet* 1965; **120**: 92–8.
- Menguy R, Desbaillets L. Role of inhibition of gastric mucous secretion in the phenomenon of gastric mucosal injury by indomethacin. *American Journal of Digestive Diseases* 1967; **12**: 862–6.
- Sarosiek J, Mizuta K, Slomiany B. Effect of acetylsalicylic acid on gastric mucin activity, permeability to hydrogen ion, and susceptibility to pepsin. *Biochem Pharmacol* 1986; **35**: 4291–5.
- Rees WDW, Bribbons LC, Turnberg LA. Effect of non-steroidal anti-inflammatory drugs and prostaglandins on alkaline secretion by rabbit gastric fundus in vitro. *Gut* 1983; **24**: 784–9.
- Baskin W, Ivey KJ, Kraus WJ, Jeffrey G, Gemmell RT. Aspirin induced ultrastructural changes in human gastric mucosa: correlation with potential differences. *Ann Intern Med* 1976; **85**: 299–303.
- Rainsford KD, Willis C. Relationship of gastric mucosal damage induced in pigs by anti-inflammatory drugs to their effect on prostaglandin production. *Dig Dis Sci* 1976; **21**: 155–64.
- Cooke AR. The role of the mucosal barrier in drugs induced gastric ulceration and erosions. *Dig Dis Sci* 1976; **21**: 155–64.
- Sarosiek J, Slomiany A, Slomiany BL. Evidence for weakening of gastric mucus integrity by *Campylobacter pylori*. *Scand J Gastroenterol* 1988; **23**: 585–90.
- Taha AS, McLaughlin S, Holland PJ, Kelly RW, Sturrock RD, Russell RI. Effect on gastric and duodenal mucosal prostaglandins of repeated intake of therapeutic doses of naproxen and etodolac in rheumatoid arthritis. *Ann Rheum Dis* 1990; **49**: 354–8.
- Redfern JS, Lee E, Feldman M. Effects of indomethacin on gastric mucosal prostaglandins in humans. Correlation with mucosal damage. *Gastroenterology* 1987; **92**: 969–77.
- Morley J, Bray MA, Jones RW, Nugteren DM, Van Drop DA. Prostaglandin and thromboxane production by human and guinea pig macrophages and leucocytes. *Prostaglandins* 1979; **17**: 730–6.
- Hawkey CJ. Synthesis of prostaglandin E₂, thromboxane B₂, and prostaglandin catabolism in gastritis and gastric ulcer. *Gut* 1986; **27**: 1484–92.
- Taha AS, Boothman P, Holland P, McKinlay A, Upadhyay R, Kelly RW, Lee F, Russell RI. Gastric mucosal prostaglandin synthesis in the presence of *Campylobacter pylori* in patients with gastric ulcer and non-ulcer dyspepsia. *Am J Gastroenterol* 1990; **85**: 47–50.
- Cryer B, Faust TW, Goldschmidt M, Redfern JS, Lee E, Feldman M. Gastric and duodenal mucosal prostaglandin concentrations in gastric or duodenal ulcer disease: relationships with demographic, environmental and histological factors. *Am J Gastroenterol* 1992; **87**: 1747–55.
- Sievert W, Lambert JR, Peacock T, Nicholson L. Gastroduodenal mucosal prostaglandin synthesis: is there an association with *Helicobacter pylori* and chronic active gastritis? *Gastroenterology* 1991; **100**: A161.
- Goren A, Fotherby KJ, Shorthouse M, Wight DGD, Hunter JO. *Campylobacter pylori* and acid secretion. *Lancet* 1989 **ii**: 212–3.
- Taha AS, Kelly RW, Gemmell CG, Lee FD, Russell RI. The interaction between *Helicobacter pylori* culture filtrate and indomethacin: effects on the integrity of human gastric antral mucosa and its prostaglandin E₂ production in vitro. *Aliment Pharmacol Therap* 1990; **4**: 265–74.
- Gana TJ, MacPherson BR, Koo J. Gastric mucosal blood flow in misoprostol pretreated aspirin-induced ulceration. *Ann Surg* 1988; **207**: 327–34.
- Arakawa T, Nakamura H, Chono S, Satoh H, Fukuda T, Saeki Y, Kobayashi K. Absence of effect of 16, 16-dimethyl prostaglandin E₂ on reduction of gastric mucosal blood flow caused by indomethacin in rats. *Dig Dis Sci* 1989; **34**: 1369–73.
- Taha AS, Angerson W, Nakshabendi J, Beekman H, Morran C, Sturrock RD, et al. Gastric and duodenal mucosal blood flow in patients receiving non-steroidal anti-inflammatory drugs – influence of age, smoking, ulceration, and *Helicobacter pylori*. *Aliment Pharmacol Therap* 1993; **7**: 41–5.
- Taha AS, Nakshabendi I, Lee FD, Sturrock RD, Russell RI. Chemical gastritis and *Helicobacter pylori* related gastritis in patients receiving non-steroidal anti-inflammatory drugs: comparison and correlation with peptic ulceration. *J Clin Pathol* 1992; **45**: 135–9.
- Rosam AC, Wallace JL, Whittle BJR. Potent ulcerogenic actions of platelet-activating factor on the stomach. *Nature* 1986; **319**: 54–6.
- Denizot Y, Shobhau J, Rambaud JC, Lewin M, Thomas Y, Benveniste J. PAF-acether synthesis by *H pylori*. *Gut* 1990; **38**: 1242–6.
- Wallace JL, Keenan CM, Granger N. Gastric ulceration induced by non-steroidal anti-inflammatory drugs is a neutrophil-dependent process. *Am J Physiol* 1990; **259**: G462–7.
- Trevethick MA, Clayton NM, Strong P, Harman IW. Do infiltrating neutrophils contribute to the pathogenesis of indomethacin induced ulceration of the rat gastric antrum? *Gut* 1993; **34**: 156–60.
- Sobala GM, King RFG, Axon ATR, Dixon MF. Reflux gastritis in the intact stomach. *J Clin Pathol* 1990; **43**: 303–6.
- Dixon MF, O'Connor HJ, Axon ATR, King RFJG, Johnston D. Reflux gastritis: distinct histopathological entity? *J Clin Pathol* 1986; **39**: 524–30.
- Laine L, Marin-Sorensen M, Weinstein WM. The histology of gastric erosions in patients taking non-steroidal anti-inflammatory drugs (NSAIDs): a prospective study. *Gastroenterology* 1988; **94**: A247.
- Graham DY, Smith JL, Spjut HJ, Torres E. Gastric adaptation: studies in humans during continuous aspirin administration. *Gastroenterology* 1988; **95**: 327–33.
- Martin DF, Montgomery E, Dobek AS, Patrissi GA, Peura DA. *Campylobacter pylori*, NSAIDs, and smoking: risk factors for peptic ulcer disease. *Am J Gastroenterol* 1989; **84**: 1268–72.
- Hudson N, Taha AS, Sturrock RD, Russell RI, Hawkey CJ. The influence of *Helicobacter pylori* colonisation on gastroduodenal ulceration in patients on non-steroidal anti-inflammatory drugs *Gut* 1992; **33**: (suppl 1) S42.
- Graham DY, Lidsky MD, Cox AM, et al. Long-term non-steroidal anti-inflammatory drug use and *Helicobacter pylori* infection. *Gastroenterology* 1991; **100**: 1653–7.
- Shallcross TM, Rathbone BJ, Wyatt JI, Heatley RV. *Helicobacter pylori*

- associated chronic gastritis and peptic ulcerations in patients taking non-steroidal anti-inflammatory drugs. *Aliment Pharmacol Therap* 1990; 4: 515-22.
- 54 Lanza FL, Evans DG, Graham DY. Effect of Helicobacter pylori infection on the severity of gastroduodenal mucosa injury after the acute administration of naproxen or aspirin to normal volunteers. *Am J Gastroenterol* 1991; 86: 735-7.
- 55 Loeb DS, Talley NJ, Ahlquist DA, Carpenter HA, Zinsmeister AR. Long-term non-steroidal anti-inflammatory drug use and gastroduodenal injury: the role of Helicobacter pylori. *Gastroenterology* 1992; 102: 1899-905.
- 56 Dixon MF, Sobala GM. Gastritis and duodenitis: the histopathological spectrum *Eur J Gastroenterol Hepatol* 1992; 4: (suppl 2) S17-23.
- 57 Taha AS, Boothman P, Nakshabendi I, Reid J, Morran C, Gemmell CG, et al. Diagnostic tests for Helicobacter pylori - comparison and influence of non-steroidal anti-inflammatory drugs. *J Clin Pathol* 1992; 45: 135-9.
- 58 Taha AS, Reid J, Boothman P, Gemmell CG, Lee FD, Sturrock RD, et al. Serological diagnosis of Helicobacter pylori - evaluation of four tests in the presence or absence of non-steroidal anti-inflammatory drugs. *Gut* 1993; 34: 461-5.
- 59 Sobala GM, Crabtree JE, Pentith JA, Rathbone JA, Shallcross TM, Wyatt JI, et al. Screening dyspepsia by serology to Helicobacter pylori. *Lancet* 1991; ii: 94-6.
- 60 Graham DY. Helicobacter pylori and the endoscopist: Whether to diagnose *Gastrointest Endosc* 1991; 37: 577-9.
- 61 Skander MP, Ryan FP. Non-steroidal anti-inflammatory drugs and pain-free peptic ulceration in the elderly. *BMJ* 1988; 297: 833-4.
- 62 Taha AS, Capell HA, Sturrock RD, Russell RI. Non-steroidal peptic damage in rheumatoid patients receiving second-line drugs. *Am J Gastroenterol* 1991; 86: 1588-91.
- 63 Caselli M, Pazzi P, La Corte R, Aleotti A, Trevisani L, Stabellini G. Campylobacter-like organisms, non-steroidal anti-inflammatory drugs and gastric lesions in patients with rheumatoid arthritis. *Digestion* 1989; 44: 101-4.
- 64 Laine L, Marin-Sorensen M, Weinstein WM. Helicobacter pylori (HP) prevalence and mucosal injury in gastric ulcers (GUs): relationship to chronic non-steroidal anti-inflammatory drug (NSAID) ingestion. *Gastroenterology* 1991; 100: A103.
- 65 Marshall BJ, McGeckie DB, Rogers PA, Glancy RJ. Pyloric campylobacter infections and gastroduodenal disease. *Med J Aust* 1985; 142: 439-44.
- 66 Von Wulffen H, Grote HJ, Gatermann S, Loning T, Berger B, Buhl C. Immunoblot analysis of immune response to Campylobacter pylori and its clinical associates. *J Clin Pathol* 1988; 41: 653-9.
- 67 Newell DG. Identification of the outer membrane proteins of Campylobacter pyloridis and antigenic cross reactivity between *C pyloridis* and *C jejuni*. *J Clin Microbiol* 1987; 133: 163-70.
- 68 Apel I, Jacobs E, Kist M, Bredt W. Antibody response of patients against a 120 kDa surface protein of Campylobacter pylori. *Zentralbl Bacteriol Microbiol Hyg [A]* 1988; 268: 271-6.
- 69 Crabtree JE, Mahony MJ, Taylor JD, Heatley RV, Littlewood JM, Tompkins DS. Immune responses to Helicobacter pylori in children with recurrent abdominal pain. *J Clin Pathol* 1991; 44: 768-71.
- 70 Crabtree JE, Taylor JD, Wyatt JI, Heatley RV, Shallcross TM, Tompkins DS, Rathbone BJ. Mucosal IgA recognition of Helicobacter pylori 120 kDa protein, peptic ulcerations and gastric pathology. *Lancet* 1991; ii: 332-5.
- 71 Figura N, Guglielmetti P, Rossolini A, Barberi A, Cusi G, Musmanno RA, Russi M, Quaranta S. Cytotoxin production by Campylobacter pylori strains isolated from patients with peptic ulcers and from patients with chronic gastritis only. *J Clin Microbiol* 1989; 27: 225-6.
- 72 Konturek SJ, Bilski J, Kwieciencin N, et al. De-Nol Stimulates gastric and duodenal alkaline secretion through prostaglandin-dependent mechanism. *Gut* 1987; 28: 1557-63.
- 73 Konturek SJ, Kwieciencin N, Obtulowicz W, et al. Effect of colloidal bismuth subcitrate on aspirin-induced gastric microbleeding, DNA loss, and prostaglandin formation in humans. *Scand J Gastroenterol* 1988; 23: 861-6.
- 74 Kent TH, Cardelli RM, Stanler FW. Small intestinal ulcers and intestinal flora in rats given indomethacin. *Am J Pathol* 1969; 54: 237-49.
- 75 Robert A, Asano T. Resistance of germ-free rats to indomethacin-induced intestinal lesions. *Prostaglandins* 1977; 14: 333-41.
- 76 Bjarnason I, Hayllar J, Smethurst P, Price A, Gumpel MJ. Metronidazole reduces intestinal inflammation and blood loss in non-steroidal anti-inflammatory drug induced enteropathy. *Gut* 1992; 33: 1204-8.
- 77 Baum C, Kennedy DL, Forbes MB. Utilization of non-steroidal anti-inflammatory drugs. *Arthritis Rheum* 1985; 28: 686-2.
- 78 Walt R, Katchinski B, Logan R, Ashley L, Langman MJS. Rising frequency of ulcer perforations in elderly people in the United Kingdom. *Lancet* 1986; i: 489-92.
- 79 Graham DY, Klein PD, Opekun AR, Boutton TW. Effect of age on the frequency of active Campylobacter pylori infection diagnosed by the [13C] urea breath test in normal subjects and patients with peptic ulcer disease. *J Infect Dis* 1988; 157: 777-80.