

APSDE-COVID statements: recommendations should be modified according to the prevalence of COVID infection rates

We read the article from Chiu *et al*¹ and would like to congratulate the authors and Asian Pacific Society for producing APSDE (Asian-Pacific Society for Digestive Endoscopy)-COVID statements. The statements make perfect practical sense in the times where robust evidence is lacking. However, there are some conflicting messages coming out and I would like to draw attention toward it.

Figure 1 is suggesting a very extensive algorithm of clinical and laboratory testing to identify the infective potential of the patient and based on that to decide if urgent or semiurgent or elective endoscopy should be performed and the type of personal protective equipment (PPE) to be used. However, statement 2 clearly indicates and quiet rightly so that in the current pandemic we should only perform endoscopy for urgent indications. I feel that the algorithm in figure 1 is best suited for countries with low incidence or recovering from pandemic and statement 2 is best suited for countries with high incidence in the midst of pandemic (the USA, Europe). A clarification along those lines will reduce the confusion and save resources and time.

The logic behind the testing for COVID-19 in patients who fulfil the criterion for urgent endoscopy is questionable. Endoscopy in these individuals is likely to be life-saving so has to be performed urgently and not delayed till test results come back. It makes no sense to take prolonged histories at least now at this stage in Europe and the USA as plenty of patients have been detected with COVID-19 infection without any contact or travel history and asymptomatic patients are infective during

endoscopy and PPE decisions on laboratory testing makes no sense either in these individuals (urgent indications for

endoscopy) as the sensitivity of test is not 100%. Recent study by Tao *et al*² reported a sensitivity rate of around 70% suggesting that a lot of these individuals with COVID-19 infection could test negative on reverse transcription PCR assays.

The best strategy for countries with high incidence rates like Europe and the USA would be to perform endoscopy for only urgent indications and all these patients should be considered as probable COVID-19 cases. When performing endoscopy for urgent indications, enhanced PPE should be the norm irrespective of clinical history and test results in high-incidence countries.

However, if we are going to perform endoscopy in known patients with COVID-19, then an additional measure of negative pressure room should be considered on top of the enhanced PPE.

The algorithm in figure 1 should be applicable when we are coming out of the pandemic or in very low incidence countries.

I would also like the authors to comment on the role of endotracheal intubation during endoscopy. I believe that the infective risks of colonoscopy is low and equally, very few colonoscopies will meet the urgent endoscopy criterion so endotracheal intubation cannot be justified. However, Endoscopic Retrograde Cholangio Pancreatography (ERCP) and gastroscopy carry a significant risk of aerosol generation and as we will only be performing these procedures for urgent indications (low numbers), so why not perform them under general anaesthesia with intubation as it will create a closed circuit with minimal risk of aerosol escaping.

Would the authors consider the above suggestions and issue an addendum in favour or against these suggestions so that the community especially in the west could be guided appropriately?

Pradeep Bhandari ^{1,2}, **Asma Alkandari**^{1,3}

¹Department of Gastroenterology, Queen Alexandra Hospital, Portsmouth, UK

²Faculty of Science, University of Portsmouth, Portsmouth, UK

³Gastroenterology, Al-Zahra Hospital, Al Jahra, Kuwait

Correspondence to Professor Pradeep Bhandari, Department of Gastroenterology, Queen Alexandra Hospital, Portsmouth PO6 3LY, UK; deep3570@yahoo.co.uk

Contributors Both authors have contributed equally to constructing and writing the contents of this letter.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, conduct, reporting or dissemination plans of this research.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; internally peer reviewed.

This article is made freely available for use in accordance with BMJ's website terms and conditions for the duration of the covid-19 pandemic or until otherwise determined by BMJ. You may use, download and print the article for any lawful, non-commercial purpose (including text and



data mining) provided that all copyright notices and trade marks are retained.

© Author(s) (or their employer(s)) 2020. No commercial re-use. See rights and permissions. Published by BMJ.

To cite Bhandari P, Alkandari A. *Gut* 2020;**69**:1367–1368.

Received 7 April 2020

Revised 13 April 2020

Accepted 14 April 2020

Published Online First 30 April 2020

Gut 2020;**69**:1367–1368. doi:10.1136/gutjnl-2020-321360

ORCID iD

Pradeep Bhandari <http://orcid.org/0000-0002-1241-2083>

REFERENCES

- 1 Chiu PWY, Ng SC, Inoue H, *et al*. Practice of endoscopy during COVID-19 pandemic: position statements of the Asian Pacific Society for digestive endoscopy (APSDE-COVID statements). *Gut* 2020;**69**:991–6.
- 2 Ai T, Yang Z, Hou H, *et al*. Correlation of chest CT and RT-PCR testing in coronavirus disease 2019 (COVID-19) in China: a report of 1014 cases. Available: <https://doi.org/10.1148/radiol.2020200642>