

## Do proton pump inhibitors influence SARS-CoV-2 related outcomes? A meta-analysis

The article by Lee *et al*<sup>1</sup> showed that the current use of proton pump inhibitors

(PPIs) increased the risk of severe clinical outcomes of COVID-19 rather than the susceptibility to SARS-CoV-2 infection in a Korean nationwide cohort. Instead, a significant association between susceptibility to SARS-CoV-2 infection and current use of PPIs, either one time or two times a day, was found by another recent study<sup>2</sup> based on US nationwide


data. The conflicting results of these two large-scale observational studies may be due to regional epidemiological differences or considerable between-study variance and might compromise clinical decision-making. As the impact of PPI use on SARS-CoV-2 infection has very relevant clinical implications, we performed a meta-analysis to address

the aforementioned discrepancies, which could lead to better informed clinical decision-making on PPI use during the ongoing pandemic.

We scrutinised 3413 records retrieved from a comprehensive search using the COVID-19 Research Articles Downloadable Database maintained by the US CDC (<https://www.cdc.gov/library/researchguides/2019novelcoronavirus/researcharticles.html>) and ultimately included 16 studies<sup>1-16</sup> from 10 countries or regions reporting comparative data on PPI use and clinical outcomes of COVID-19 (online supplemental figure 1 and table). We pooled the data using an inverse variance-weighted random-effect model. Pooled estimates are presented as OR, HR or mean difference (MD), with associated 95% CIs. Intensive care unit admission, mechanical ventilation, acute respiratory distress syndrome or death were considered severe outcomes of COVID-19.


Six studies<sup>1-6</sup> including 318 261 participants reported data on PPI usage and the risk of SARS-CoV-2 infection. Among them, five studies had information of current PPI users compared with non-users and four on past PPI users versus non-users. Analysis of five studies<sup>1-5</sup> encompassing 145 428 patients who were tested for SARS-CoV-2 showed that the risk of SARS-CoV-2 infection was higher, although not significantly, among current PPI users (OR 1.33, 95% CI 0.86 to 2.07,  $p=0.20$ ; figure 1) compared with PPI non-users, with evidence of substantial between-study heterogeneity ( $I^2=97\%$ ). Moreover, in a subgroup analysis of non-Korean cohorts,<sup>2-4</sup> we found a significant association between current use of PPIs and increased risk of SARS-CoV-2 infection (OR 1.94, 95% CI 1.59 to 2.36,  $p<0.0001$ ; online supplemental figure 2). Furthermore, a leave-one-out sensitivity analysis revealed that the summary estimate of the association between current PPI usage and SARS-CoV-2 infection was overly influenced by a single Korean study<sup>5</sup> (online supplemental figure 3).

Instead, current or regular PPI users were more likely to have severe outcomes of COVID-19 than PPI non-users, with a pooled OR of 1.67 (95% CI 1.19 to 2.33,  $p=0.003$ ;  $n=42\,405$  from nine studies;<sup>1 3 7-13</sup>  $I^2=63\%$ ; figure 2) and a pooled HR of 1.87 (95% CI 1.29 to 2.70,  $p<0.001$ ;  $n=2977$  from two studies;<sup>15 16</sup>  $I^2=80\%$ ; figure 2). These results were consistent with our leave-one-out sensitivity analysis (online supplemental figure 4), indicating that this association was strong. Furthermore,




**Figure 1** Forest plot showing the association between PPI use and SARS-CoV-2 infection. PPI, proton pump inhibitor.


#### A Severe outcomes of COVID-19 (expressed as Odds Ratio)



#### B Severe outcomes of COVID-19 (expressed as Hazard Ratio)



#### C Duration of hospital stay



**Figure 2** Forest plot showing the association of PPI use with severe outcomes of COVID-19 (A, OR; B, HR) or duration of hospital stay (C). PPI, proton pump inhibitor.

current PPI users tended to hospitalised longer than PPI non-users, although not by a statistically significant margin ( $n=353$  from two studies;<sup>7,14</sup> MD 1.13, 95% CI  $-0.18$  to  $2.43$ ,  $p=0.09$ ; figure 2). Finally, past use of PPIs was not associated with increased susceptibility to SARS-CoV-2 infection ( $n=172833$  from four studies;<sup>13,16</sup> OR 0.85, 95% CI 0.57 to 1.27,  $p=0.43$ ;  $I^2=92\%$ ; figure 1) or with severe outcomes of COVID-19 ( $n=40097$  from three studies;<sup>13,9</sup> OR 1.03, 95% CI 0.85 to 1.23,  $p=0.79$ ;  $I^2=0\%$ ; figure 2).

In summary, this meta-analysis shows that regional differences can explain the heterogeneous findings concerning the association between current PPI use and incidence of SARS-CoV-2 infection and further underscores the increased risk of severe COVID-19 outcomes associated with current PPI use, highlighting that caution should be exercised when treating patients receiving PPIs during the COVID-19 pandemic. Further studies investigating different dosing regimens and durations of PPI use on COVID-19 outcomes should be warranted.

Guo-Fu Li<sup>1,2</sup>, Xiao-Xiao An,<sup>2,3</sup> Yichao Yu,<sup>4</sup> Li-Rong Jiao,<sup>2,3</sup> Daniele Canarutto,<sup>5</sup> Guo Yu<sup>1,2</sup>, Guangji Wang,<sup>6</sup> Dan-Na Wu,<sup>7</sup> Yin Xiao<sup>8</sup>

<sup>1</sup>Clinical Medical College, Yangzhou University, Yangzhou, China

<sup>2</sup>Institution of Drug Clinical Trial, Subei People's Hospital, Yangzhou, China

<sup>3</sup>College of Pharmacy, Dalian Medical University, Dalian, Liaoning, China

<sup>4</sup>Department of Pharmaceutics, University of Florida, Gainesville, Florida, USA

<sup>5</sup>Faculty of Medicine and Surgery, Vita Salute San Raffaele University, Milan, Italy

<sup>6</sup>Key Laboratory of Drug Metabolism and Pharmacokinetics, China Pharmaceutical University, Nanjing, China

<sup>7</sup>Department of Pharmacy, Hainan General Hospital (Hainan Affiliated Hospital of Hainan Medical University), Haikou, China

<sup>8</sup>Department of Pharmacy, Haikou Affiliated Hospital of Central South University Xiangya School of Medicine, Haikou, China

**Correspondence to** Dr Guo Yu, Clinical Medical College, Yangzhou University, Yangzhou 225009, China; guoyu@yzu.edu.cn

**Contributors** Concept and design: G-FL and GY. Acquisition, analysis and interpretation of data: G-FL, X-XA, GY, YY, L-RJ, D-NW, YX. Drafting of the

manuscript: GFL. Supervision: GY. Critical revision of the manuscript: DC, G-FL, GW and YY. Final approval: all authors.

**Funding** This work was supported by Jiangsu Provincial Medical Youth Talent programme (QNRC2016323), Jiangsu Province 333 Project (to GY) and Jiangsu Provincial Science Fund for Distinguished Young Scholars (to GY).

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Provenance and peer review** Not commissioned; externally peer reviewed.

This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.



## OPEN ACCESS

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

CC-BY-NC

► Additional material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/gutjnl-2020-323366>).



**To cite** Li G-F, An X-X, Yu Y, et al. *Gut* 2021;**70**:1806–1808.

Received 13 October 2020

Revised 28 October 2020

Accepted 30 October 2020

Published Online First 10 November 2020

*Gut* 2021;**70**:1806–1808. doi:10.1136/gutjnl-2020-323366

**ORCID iDs**

Guo-Fu Li <http://orcid.org/0000-0002-4628-9941>


Guo Yu <http://orcid.org/0000-0001-6685-2167>

## REFERENCES

- Lee SW, Ha EK, Yeniova Abdullah Özgür. Severe clinical outcomes of COVID-19 associated with proton pump inhibitors: a nationwide cohort study with propensity score matching. *Gut* 2021;**70**:807–9.
- Almaro CV, Chey WD, Spiegel BMR, et al. Increased risk of COVID-19 among users of proton pump inhibitors. *Am J Gastroenterol* 2020;**115**:1707–15.
- Ullah A, Sivapalan L, Chelala PC, et al. COVID-19 in patients with hepatobiliary and pancreatic diseases in East London: a single-centre cohort study. *medRxiv* 2020.
- Vila-Corcoles A, Satue-Gracia E, Ochoa-Gondar O, et al. Use of distinct anti-hypertensive drugs and risk for COVID-19 among hypertensive people: a population-based cohort study in southern Catalonia, Spain. *J Clin Hypertens* 2020. doi:10.1111/jch.13948. [Epub ahead of print: 25 Jul 2020].
- Huh K, Ji W, Kang M, et al. Association of previous medications with the risk of COVID-19: a nationwide claims-based study from South Korea. *medRxiv* 2020.
- Tarlow B, Gubatan J, Khan MA, et al. Are proton pump inhibitors contributing to SARS-COV-2 infection? *Am J Gastroenterol* 2020. doi:10.14309/ajg.0000000000000933. [Epub ahead of print: 11 Sep 2020].
- Ramachandran P, Perisetti A, Gajendran M, et al. Prehospitalization proton pump inhibitor (PPI) use and clinical outcomes in COVID-19. *medRxiv* 2020.
- Luxemburger H, Sturm L, Biever P, et al. Treatment with proton pump inhibitors increases the risk of secondary infections and ARDS in hospitalized patients with COVID-19: coincidence or underestimated risk factor? *J Intern Med* 2020. doi:10.1111/joim.13121. [Epub ahead of print: 01 Jul 2020].
- McKeigue PM, Kennedy S, Weir A, et al. Associations of severe COVID-19 with polypharmacy in the REACT-SCOT case-control study. *medRxiv* 2020.
- Argenziano MG, Bruce SL, Slater CL, et al. Characterization and clinical course of 1000 patients with coronavirus disease 2019 in New York: retrospective case series. *BMJ* 2020;**369**:m1996.
- Cheung KS, Hung IFN, Leung WK. Association between famotidine use and COVID-19 severity in Hong Kong: a territory-wide study. *Gastroenterology* 2020.
- Lossner M-R, Lapoix C, Delannoy M, et al. Almitrine as a non-ventilatory strategy to improve intrapulmonary shunt in COVID-19 patients. *Anaesth Crit Care Pain Med* 2020;**39**:467–9.
- Yan S, Song X, Lin F, et al. Clinical characteristics of coronavirus disease 2019 in Hainan, China. *medRxiv* 2020.
- Zhang XY, HB W, Ling Y, et al. Analysis of the effect of proton pump inhibitors on the course of common COVID-19. *MedRxiv* 2020.
- Jimenez L, Codo AC, Sampaio VS, et al. The influence of pH on SARS-CoV-2 infection and COVID-19 severity. *MedRxiv* 2020.
- Freedberg DE, Conigliaro J, Wang TC, et al. Famotidine use is associated with improved clinical outcomes in hospitalized COVID-19 patients: a propensity score matched retrospective cohort study. *Gastroenterology* 2020;**159**:1129–31.


**Supplementary figure 1: Flow chart for study selection**

\*Searches using strategy#1 ("proton pump inhibitor\*" or "PPI\*" OR " H2-receptor antagonist\*" OR hypochlorhydria OR "gastric acid" OR "gastric pH" OR omeprazole OR rabeprazole OR esomeprazole OR famotidine OR pantoprazole OR lansoprazole) or strategy#2 (gastrointestinal[title/abstract]) were performed in the COVID-19 Research Articles Downloadable Database by the US CDC (<https://www.cdc.gov/library/researchguides/2019novelcoronavirus/researcharticles.html>), which includes literature from 25 databases, such as Medline (Ovid and PubMed), Embase, Scopus, Cochrane Library, LitCovid, WHO COVID-19 website, medRxiv (preprints), bioRxiv (preprints), chemRxiv (preprints), and SSRN (preprints).




Supplementary figure 1


Supplementary figure 2: Subgroup analysis of Korean versus non-Korean cohorts for the association between PPI use and risk of SARS-CoV-2 infection



### Supplementary figure 3: Forest plot showing leave-one-out sensitivity analysis for the association of PPI use with incidence of SARS-CoV-2 infection



### Supplementary figure 4: Forest plot showing leave-one-out sensitivity analysis for the association of PPI use with severe outcomes of COVID-19





Supplementary table: Summary characteristics of the included studies

Study	Study design	Country or region	Timing of data collection	Mean or median age (years)	Male subjects (%)	Number of subjects	Number of PPI users	Clinical outcome	Confounder adjustment
Lee 2020 <sup>1</sup>	Retrospective cohort	Korea	Jan 1 to May 15, 2020	48	51.0	132316	20405	SARS-CoV-2 infection; severe outcomes of COVID-19*	Yes
Almario 2020 <sup>2</sup>	Retrospective cohort	USA	May 3 to Jun 24, 2020	NR	48	53130	16547	SARS-CoV-2 infection	Yes
Ullah 2020 <sup>3</sup>	Retrospective cohort	UK	Feb 12 to Jun 12, 2020	57	43.9	15586	5908	SARS-CoV-2 infection; severe outcomes of COVID-19*	No
Corcoles 2020 <sup>4</sup>	Retrospective cohort	Spain	May 1 to Apr 3, 2020	≥ 50	48.1	34936	11807	SARS-CoV-2 infection	No
Huh 2020 <sup>5</sup>	Case-control	Korea	Up to Apr 8, 2020	49	48.7	65149	14167	SARS-CoV-2 infection	Yes
Tarlow 2020 <sup>6</sup>	Retrospective cohort	USA	NR	NR	NR	84325	18240	SARS-CoV-2 infection	No
Ramachandran 2020 <sup>7</sup>	Retrospective cohort	USA	Mar 1 to Apr 25, 2020	66	54.9	295	46	Severe outcomes of COVID-19*; duration of hospital stay	Yes
Luxenburger 2020 <sup>8</sup>	Retrospective cohort	Germany	NR	65	56.6	152	62	Severe outcomes of COVID-19*	No
McKeigue 2020 <sup>9</sup>	Case-control	Scotland	Up to Jun 6, 2020	NR	NR	41220	2715	Severe outcomes of COVID-19*	No
Argenziano 2020 <sup>10</sup>	Retrospective cohort	USA	Mar 1 to Apr 5, 2020	63	59.6	1000	163	Severe outcomes of COVID-19*	No
Cheung 2020 <sup>11</sup>	Retrospective cohort	Hongkong	Jan 1 to May 10, 2020	NR	NR	952	27	Severe outcomes of COVID-19*	Yes
Losser 2020 <sup>12</sup>	Case series (individual)	France	Mar16 to Apr 12, 2020	70	58.8	17	6	Severe outcomes of COVID-19*	No
Yan 2020 <sup>13</sup>	Retrospective cohort	China	Jan 22 to Mar 13, 2020	51	48.2	168	32	Severe outcomes of COVID-19*	No
Zhang 2020 <sup>14</sup>	Retrospective cohort	China	Jan 20 to Mar 16, 2020	50	55.2	58	29	Duration of hospital stay	Yes



Jimenez 2020 <sup>15</sup>	Retrospective cohort	Brazil	NR	NR	NR	1357	242	Severe outcomes of COVID-19*	Yes
Freedberg 2020 <sup>16</sup>	Retrospective cohort	USA	Feb 25 to Apr 13, 2020	NR	NR	1620	NR	Severe outcomes of COVID-19*	Yes

\*Severe outcomes of COVID-19 consisted of admission to the intensive care unit, mechanical ventilation, acute respiratory distress syndrome, or death.  
COVID-19, Coronavirus Disease 2019; NR, not reported; PPI, proton pump inhibitor; SARS-CoV- 2, severe acute respiratory syndrome coronavirus 2.