

## Materials and Methods

**Study cohort and approval.** The specific use of each autopsy sample in each experiment is provided in [supplementary table 1](#). All safety precautions were in line with recently published guidelines.[1] The study of autopsy samples does not meet the definition of Human Subject Research in Singapore, and all samples were anonymously coded in accordance with the Helsinki Declaration. The Agency of Science, Technology and Research, Singapore, provided approval for the use of control tissue materials in this study IRB: 2020 112.

**Digital spatial profiling.** NanoString spatial profiling[2, 3] was used to analyze 3 protein-level and 1,800 transcriptomic-level immune-markers simultaneously on formalin-fixed paraffin-embedded (FFPE) tissue slides. Following deparaffinization and antigen retrieval procedures, sections were simultaneously incubated overnight with fluorescent-labeled antibodies against CD3 (Origene), CD68 (Santa Cruz), and Pan-Cytokeratin (Novus Biologicals) or cytokeratin 8/18 (Novus Biologicals) to visualize morphological features in the regions of interest (ROIs). After staining, the tissues were scanned using a GeoMx DSP instrument to generate digital fluorescent images and select individual ROIs inside a geometric section. To carry out high-resolution multiplex profiling, each ROI was assigned as a CD3<sup>+</sup> T-cell-rich or CD68<sup>+</sup> macrophage-rich region. UV light was directed through a programmable digital micromirror device (DMD) or dual DMD (DDMD) to accurately illumine the ROI and cleave the photocleavable oligos (PC-oligos) from the selected region, which were then collected by microcapillary tube inspiration and dispensed into a 96-well plate. Inside the microplate, the single-molecule counting nCounter System enabled the digital counting of released oligos.[4, 5] Individual counts were normalized against the 75th percentile of the signal from their own ROI (Q3 normalization).[6]

**Multiplex immunohistochemistry.** Multiplex immunohistochemistry was performed using an Opal Multiplex IHC kit (Akoya Biosciences, California), as previously described.[7, 8, 9, 10] In brief, FFPE tissue sections were cut onto Bond Plus slides (Leica Biosystems, Richmond) and heated at 60°C for 20 min.[11] The tissue slides were subjected to deparaffinization, rehydration, and heat-induced epitope retrieval using a Leica Bond Max autostainer (Leica Biosystems, Melbourne) before endogenous peroxidase blocking (Leica Biosystems, Newcastle). Next, the slides were incubated with primary antibodies followed by incubation with polymeric HRP-conjugated secondary antibodies (Leica Biosystems, Newcastle) (supplementary table 2). The samples were incubated with Opal fluorophore-conjugated tyramide signal amplification (TSA) (Akoya Biosciences, California) at 1:100 dilution. The slides were rinsed with wash buffer (BOND Wash Solution 10X Concentrate) after each step. Following TSA deposition, the slides were again subjected to heat-induced epitope retrieval to strip the tissue-bound primary/secondary antibody complexes before further labeling. These steps were repeated until the samples were labeled with all six markers and spectral DAPI (Akoya Biosciences, California) at a 1:10 dilution. Finally, the slides were mounted in ProLong Diamond Anti-fade Mountant (Molecular Probes, Life Technologies, USA) and developed in the dark at room temperature for 24 h. Images were captured for each case under a Vectra 3 pathology imaging system microscope (Akoya Biosciences, California) and then analyzed and scored by a pathologist using inForm software (version 2.4.2; Akoya Biosciences) and HALO™ (Indica Lab). Raw images have been deposited in <https://immunoatlas.org/MIHC/210723-2/MIHC21048/> and <https://immunoatlas.org/MIHC/210723-2/MIHC21049/>.

**Supplementary figure 1: Biological pathways enriched in COVID-19/HIV co-infection**

(A) Relative estimated levels of immune cells determined by deconvolution of digital spatial profiling ROIs using CIBERSORTx (<https://cibersort.stanford.edu/>). (B) Biological pathways enriched in the liver, kidney, and lung ROIs of the COVID-19/HIV decedent compared with the COVID-19 decedents identified by gene set enrichment analysis. A positive enrichment score indicates enrichment of the process in the COVID-19/HIV decedent.

**Supplementary figure 2: Increased macrophage response in the liver of the COVID-19/HIV decedent is associated with COVID-19 viral load**

(A) Expression heatmap of genes related to T-cell activation or suppression in CD3-rich ROIs (*left*) and myeloid activation in CD68-rich ROIs (*right*) of the kidney. (B) Correlation between NP<sup>+</sup> cells and T-cells in ROIs of the lung of the COVID-19/HIV (red;  $p=0.06$ ) and COVID-19 (blue;  $p=0.0004$ ) decedents.  $p$ -values were calculated by Wald tests. (C) Comparison of the abundance of activated macrophages in the liver between the COVID-19/HIV decedent and COVID-19 decedents. The  $p$ -value was calculated by a two-tailed U-test: \*(<0.05).

**Supplementary table 1: Study cohort and tissues obtained for analysis**

<b>Patient ID</b>	<b>Sex</b>	<b>Survival since symptom onset (days)</b>	<b>Digital Spatial Profiling</b>	<b>mIHC</b>	<b>Clinical Diagnosis</b>	<b>Cause(s) of Death</b>
Pt-HIV	Male	46	Lung, Liver, Kidney	Lung, Liver, Kidney	COVID-19; HIV;	RFRWS
Pt-CV1	Male	20	Lung, Liver, Kidney	Lung, Liver, Kidney	COVID-19	RFRWS
Pt-CV2	Male	21	Lung, Liver	Lung, Liver	COVID-19	RFRWS
Pt-CV3	Male	30	Lung, Liver	Lung, Liver	COVID-19	RFRWS

*Abbreviation: RFRWS: Respiratory failure related with SARS-CoV-2; mIHC: multiplex immunohistochemistry.*

**Supplementary table 2: Antibodies used for digital spatial profiling and multiplex immunohistochemistry**

Usage	Primary antibody	Company	Clone
Digital Spatial Profiling	CD3	Origene	UMAB54
	CD68	Santa Cruz	KP1
	PanCK	Novus Biologicals	AE1/AE3
	CK8/18	Novus Biologicals	KRT8/803 + KRT18/835
Multiplex immunohistochemistry	BATF3	Abcam	Polyclonal
	BDCA2	Merck	10E6.1
	CD206	Santa Cruz	D-1
	CD3	Dako	Polyclonal
	CD80	Abcam	EPR1157(2)
	CD83	BioLegend	HB15e
	SARS-CoV-2 (NP)	Novus Biologicals	Polyclonal

## References

- 1 Hanley B, Lucas SB, Youd E, Swift B, Osborn M. Autopsy in suspected COVID-19 cases. *Journal of Clinical Pathology* 2020;**73**:239.
- 2 Delorey TM, Ziegler CGK, Heimberg G, Normand R, Yang Y, Segerstolpe Å, *et al.* COVID-19 tissue atlases reveal SARS-CoV-2 pathology and cellular targets. *Nature* 2021.
- 3 Rendeiro AF, Ravichandran H, Bram Y, Chandar V, Kim J, Meydan C, *et al.* The spatial landscape of lung pathology during COVID-19 progression. *Nature* 2021;**593**:564-9.
- 4 Beechem JM. High-Plex Spatially Resolved RNA and Protein Detection Using Digital Spatial Profiling: A Technology Designed for Immuno-oncology Biomarker Discovery and Translational Research. *Methods Mol Biol* 2020;**2055**:563-83.
- 5 Toki MI, Merritt CR, Wong PF, Smithy JW, Kluger HM, Syrigos KN, *et al.* High-Plex Predictive Marker Discovery for Melanoma Immunotherapy-Treated Patients Using Digital Spatial Profiling. *Clinical cancer research : an official journal of the American Association for Cancer Research* 2019;**25**:5503-12.
- 6 Beasley GM, Therien AD, Holl EK, Al-Rohil R, Selim MA, Farrow NE, *et al.* Dissecting the immune landscape of tumor draining lymph nodes in melanoma with high-plex spatially resolved protein detection. *Cancer Immunology, Immunotherapy* 2020.
- 7 Lim JCT, Yeong JPS, Lim CJ, Ong CCH, Wong SC, Chew VSP, *et al.* An automated staining protocol for seven-colour immunofluorescence of human tissue sections for diagnostic and prognostic use. *Pathology* 2018;**50**:333-41.
- 8 Yeong J, Lim JCT, Lee B, Li H, Ong CCH, Thike AA, *et al.* Prognostic value of CD8 + PD-1+ immune infiltrates and PDCD1 gene expression in triple negative breast cancer. *Journal for ImmunoTherapy of Cancer* 2019;**7**:34.
- 9 Yeong J, Thike AA, Lim JC, Lee B, Li H, Wong SC, *et al.* Higher densities of Foxp3(+) regulatory T cells are associated with better prognosis in triple-negative breast cancer. *Breast Cancer Res Treat* 2017;**163**:21-35.
- 10 Cheung CCL, Goh D, Lim X, Tien TZ, Lim JCT, Lee JN, *et al.* Residual SARS-CoV-2 viral antigens detected in GI and hepatic tissues from five recovered patients with COVID-19. *Gut* 2021:gutjnl-2021-324280.
- 11 Yeong J, Lim JCT, Lee B, Li H, Chia N, Ong CCH, *et al.* High Densities of Tumor-Associated Plasma Cells Predict Improved Prognosis in Triple Negative Breast Cancer. *Frontiers in immunology* 2018;**9**:1209-.